

Government of the People's Republic of Bangladesh  
**Bangladesh Food Safety Authority Ministry of Food**

**DRAFT NOTIFICATION**

Dated the \_\_\_\_\_

**S.R.O. No. \_\_\_\_\_** - In exercise of the powers conferred by section 87 of the Food Safety Act 2013 (Act No. 43 of 2013), to be read with section 13 (4), the Bangladesh Food Safety Authority, with the prior approval of the Government hereby enacted the following Regulation, as follows: -

**(1) Title and commencement.** - (i) This Regulations may be called the Food Safety (Pesticides and Other Chemical Residues) Regulation, 2023.

(ii) It shall come into force after 6 month from the date of their publication in the Official Gazette and Food Business Operator shall comply with all the provisions of these regulations by -----.

**(2) Definitions.** - (i) In this Regulations, unless there is anything repugnant in the subject or context,

(ii) **“Maximum Residue Limit”** means the maximum level of a pesticide residue that is legally permitted in or on food or feed.

(iii) **“Act”** means the Food Safety Act, 2013 (Act No. 43 of 2013);

(iv) **“Pesticide”** means any substance or mixture of substances used or represented as a means for preventing, destroying, repelling, mitigating, or controlling, directly or indirectly, any pest, fungus, bacterial, organism, nematode, virus, weed, rodent, or other plant or insect; but does not include any substance which is a ‘drug’ within the meaning of the Drugs Act, 1940;

(v) **“Pesticide Residues”** means any specified substance in food or any such condition thereof that results from the use of insecticides or pesticides during its production, importation, processing, storing, supply, sale or marketing and contains the parent compound of the insecticides or pesticides, derivatives, conversion products, residue of metabolites, reaction products or created impurities and presence of which is considered to significantly intoxicate any food product, and also includes the residues contaminated from environment;

(vi) The words or expressions used in this regulation but not provided with any definition or explanation shall carry the same meaning as provided under the Act.

**(3) Controlling use of pesticide residues in articles of food.** - Any person or his/her agent, shall not produce, import, process, stock, supply, market or sell any article of food which is harmful to health for containing residue of pesticide in excess of the maximum residue limit specified in the Schedule 1.

**(4) Portion of commodities to which maximum residue limits apply and which is analyzed.** - Maximum Residue Limits stipulated in Schedule-1 and Schedule-2 are in most cases stated in terms of a specific whole raw agricultural commodity as it moves in trade. In some instances, a qualification is included that describes the part of the raw agricultural commodity to which the maximum residue limit applies, for example, almonds on a shell-free basis and beans without pods.



In other instances, such qualifications are not provided. Therefore, unless otherwise specified, the portion of the raw agricultural commodity to which the MRL applies and which is to be prepared as the analytical sample for the determination of pesticide residues is as described in the Schedule-3.

**(5) Controlling use of harmful residues in articles of food.**-Any person or his agent, shall not produce, import, process, stock, supply, market or sell any article of food which is harmful to health for containing residue of plant growth regulator specified from Schedule-2 in excess of the Maximum Residue Limits specified in the abovementioned schedules.

**(6) Following international standard.** - In order to determine the maximum residue limit of pesticide and plant growth regulator with respect to the production, processing, import, marketing or sell of any article of food not mentioned within Schedule-1 to Schedule-3 respectively, the limit as stipulated in the latest edition of Codex Alimentarius or the internationally recognized standard (as appropriate) shall have to be followed.

**(7) Propagation of advertisement.** -In order to fulfill the objective of section 41 and 42 of the Act, any person or his agent shall not propagate or sell any advertisement for the production, import, marketing or sell of any article of food containing pesticide and other chemical residues in excess of the maximum residue limit specified from Schedule-1 and Schedule-2 which may be harmful for the buyer or may mislead public at large.

**(8) Inapplicability.** -The provisions of Pure Food Rules 1967 and Food Safety (Chemical Contaminant, Toxin and Harmful Residues) Regulation, 2017 concerned with this Regulation shall be rendered inapplicable as soon as this Regulation comes into effect.

**(9) Publication of English translation.** - After the commencement of this Regulation, the Authority, with prior approval of the Government, if necessary, by notification in the official Gazette, may publish an Authentic English text of the Regulation.

Provided that, in the event of conflict between the Bangla and the English text, the Bangla text shall prevail.



**Schedule-1**  
**Maximum Residues Limits (MRL) of Pesticides**

Name of the Pesticide(s)	Name of the Food	MRL (mg/kg)	Symbol	Notes
2,4-D Amine Salt	Tea	0.05		
2,4-Dichlorophenoxy Acetic Acid (2,4-D)	Berries and other small fruits	0.1		
	Citrus fruits (group)	1	Po	
	Edible offal (mammalian)	5		
	Eggs	0.05		(Shell free basis)
	Fruits	2		
	Maize	0.05		
	Meat (from mammals other than marine mammals)	0.2		
	Milks	0.05		
	Milled food grains	0.01		
	Other food grains	0.01		
	Pome fruits (group)	0.01	(*)	
	Potato	0.2		
	Poultry meat	0.05	(*)	
	Poultry, edible offal of	0.05	(*)	
	Rice, hay and/or straw	10		
	Rice, husked	0.1		
	Rye	2		
	Sorghum grain	0.01	(*)	
	Soya bean (dry)	0.01	(*)	
	Stone fruits (group)	0.05	(*)	
	Sugar cane	0.05		
	Sweet corn (corn-on-the-cob)	0.05	(*)	
	Tree nuts (group)	0.2		
	Wheat	2		
	Wheat, hay and/or straw	100		
Abamectin	Almond hulls	0.2		
	Avocado	0.01		
	Beans (dry)	0.005		
	Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	0.08		(immature beans with pods)
	Cane berries (subgroup)	0.2		
	Celery	0.03		
	Cherries (subgroup)	0.07		
	Chive, dry	0.08		
	Citrus fruits (group)	0.02		
	Cotton seed	0.01		

	Cucumber	0.03		
	Egg plant	0.05		
	Garlic	0.005		
	Gherkin	0.03		
	Grape juice	0.05		
	Grape, dried (= Currants, Raisins and Sultanias)	0.1		
	Grapes	0.05	*	
	Green onions (subgroup)	0.01		
	Herbs	0.01		
	Hops, dry	0.15		
	Lettuce, head	0.15		
	Mango	0.01		
	Melons, except watermelon	0.01		
	Onion, bulb	0.005		
	Orange oil, edible	0.1		
	Papaya	0.01		
	Peaches (including apricots and nectarine) (subgroup)	0.03		
	Peanut	0.005	(*)	
	Chilli	0.05	*	
	Peppers chili, dried	0.5		
	Peppers, sweet (including pimento or pimienta)	0.09		
	Pineapple	0.002	(*)	
	Plums (including fresh prunes) (subgroup)	0.005		
	Pome fruits (group)	0.01		
	Potato	0.005	(*)	
	Rice, hay and/or straw	0.001		
	Rice, husked	0.002		
	Shallot	0.005		
	Soya bean (dry)	0.002	(*)	
	Strawberry	0.15		
	Succulent beans without pods (subgroup)	0.002	(*)	
	Sweet corns (subgroup)	0.002		
	Sweet potato	0.005	(*)	
	Tomato	0.05		
	Tree nuts (group)	0.005	(*)	
	Yams	0.005	(*)	
Acephate (expressed as mixture of Methamidophos & acephate)	Artichoke, globe	0.3		



	Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	5		
	Cabbages, head	2		
	Cranberry	0.5		
	Edible offal (mammalian)	0.05		
	Eggs	0.01	(*)	
	Meat (from mammals other than marine mammals)	0.05		
	Milks	0.02		
	Peppers chili, dried	50		
	Poultry fats	0.1		
	Poultry meat	0.01	(*)	
	Poultry, edible offal of	0.01	(*)	
	Rice, hay and/or straw	0.3		
	Rice, husked	1		
	Soya bean (dry)	0.3		
	Spices	0.2	(*)	
	Tomato	1		
	Safflower seed	2		
	Cotton seed	2		
Acetamiprid	Asparagus	0.8		
	Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	0.4		
	Beans, shelled	0.3		
	Berries and other small fruits	2		except grapes and strawberries
	Cabbages, head	0.7		
	Celery	1.5		

	Cherries (subgroup)	1.5		
	Citrus fruits (group)	1		
	Cotton seed	0.7		
	Cucumber	0.3		
	Edible offal (mammalian)	1		
	Eggs	0.01	(*)	
	Flowerhead brassicas (includes Broccoli: Broccoli, Chinese and cauliflower)	0.4		
	Fruiting vegetables, cucurbits (group)	0.2		(except cucumber)
	Fruiting vegetables, other than cucurbits (group)	0.2		except sweet corn & mushrooms
	Garlic	0.02		
	Grapes	0.5		
	Mammalian fats (except milk fats)	0.3		
	Meat (from mammals other than marine mammals)	0.5		
	Milks	0.2		
	Nectarine	0.7		
	Onion, bulb	0.02		
	Peach	0.7		
	Peas, shelled (succulent seeds)	0.3		
	Pepper, black, white	0.1		
	Peppers chili, dried	2		
	Plums (including fresh prunes) (subgroup)	0.2		except prunes
	Pome fruits (group)	0.8		
	Poultry meat	0.01	(*)	
	Poultry, edible offal of	0.05	(*)	
	Prunes	0.6		
	Spices, seeds	2		
	Spring onion	5		
	Strawberry	0.5		
	Sweet corn (corn-on-the- cob)	0.01		
	Tree nuts (group)	0.06		
	Okra	0.1		
	Cabbage	0.7		
	Cotton seed oil	0.1		
Acetochlor	Alfalfa hay	30		
	Barley	0.04		
	Barley, hay and/or straw	0.3		

	Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	0.02	(*)	
	Broad bean (dry)	0.15		
	Buckwheat	0.04	(*)	
	Chick-pea (dry)	0.15	(*)	
	Edible offal (mammalian)	0.05		
	Eggs	0.02	(*)	
	Hyacinth bean (dry)	0.15		
	Lentil (dry)	0.15		
	Lupin (dry)	0.15		
	Maize	0.02		
	Mammalian fats (except milk fats)	0.02	(*)	
	Meat (from mammals other than marine mammals)	0.02	(*)	
	Milks	0.02	(*)	
	Millet (Including barnyard, bulrush, common, finger, foxtail, little Millet)	0.04	(*)	
	Oats	0.04	(*)	
	Peas (dry)	0.02	(*)	
	Pigeon pea (dry)	0.15		
	Potato	0.04	(*)	
	Poultry meat	0.02	(*)	
	Poultry, edible offal of	0.02	(*)	
	Rye	0.04	(*)	
	Soya bean (dry)	1.5		
	Sugar beet	0.15		
	Sugar beet leaves or tops (dry)	3		
	Sugar beet molasses	0.3		
	Sugar beet pulp, dry	0.3		
	Sunflower seed	0.04		
	Sweet corn (corn-on-the- cob)	0.04		
	Teosinte	0.04	(*)	
	Triticale	0.04	(*)	
	Wheat	0.02	(*)	
	Wheat, hay and/or straw	0.2		
	Wild rice	0.04	(*)	
Alpha Cypermethrin	Cottonseed Oil	0.05		
Ametoctradin	Brassica vegetables (except Brassica leafy vegetables) (group)	9		
	Celery	20		
	Cucumber	0.4		



	Eggs	0.03	(*)	
	Fruiting vegetables, cucurbits (group)	3		(except cucumber)
	Fruiting vegetables, other than cucurbits (group)	1.5		(except sweet corn and mushroom)
	Garlic	1.5		
	Grape, dried (= Currants, Raisins and Sultanas)	20		
	Grapes	6		
	Hops, dry	30		
	Leafy vegetables	50		
	Onion, bulb	1.5		
	Peppers chili, dried	15		
	Potato	0.05		
	Poultry fats	0.03	(*)	
	Poultry meat	0.03	(*)	
	Poultry, edible offal of	0.03	(*)	
	Shallot	1.5		
	Spring onion	20		
Ametyrne	Sugarcane	0.05	*	
Amitraz	Cattle meat	0.05		The MRL accommodates external animal treatment.
	Cherries (subgroup)	0.5		
	Cotton seed	0.5		
	Cotton seed oil, crude	0.05		
	Cucumber	0.5		
	Edible offal of cattle, pigs & sheep	0.2		The MRL accommodates external animal treatment.
	Milks	0.01	(*)	The MRL accommodates external animal treatment.
	Oranges, sweet, sour (including Orange-like hybrids) (subgroup)	0.5		
	Peach	0.5		
	Pig meat	0.05		The MRL accommodates external animal treatment.
	Pome fruits (group)	0.5		



	Sheep meat	0.1		The MRL accommodates external animal treatment.
	Tomato	0.5		
Atrazine	Maize	0.01		
	Sugarcane	0.25		
Azocyclotin	Apple	0.2		
	Currants, black, red, white	0.1		
	Grapes	0.3		
	Oranges, sweet, sour (including Orange-like hybrids) (subgroup)	0.2		
	Pear	0.2		
Azoxystrobin	Almond hulls	7		
	Artichoke, globe	5		
	Asparagus	0.01	(*)	
	Banana	2		
	Barley	1.5		
	Berries and other small fruits	5		Except cranberry, grapes and strawberry
	Brassica vegetables (except Brassica leafy vegetables) (group)	5		
	Bulb vegetables (group)	10		
	Carambola	0.1		
	Celery	5		
	Citrus fruits (group)	15		
	Coffee beans	0.03		
	Cotton seed	0.7		
	Cranberry	0.5		
	Dried herbs	300		Except hops (dry)
	Edible offal (mammalian)	0.07		
	Eggs	0.01	(*)	
	Fruiting vegetables, cucurbits (group)	1		
	Fruiting vegetables, other than cucurbits (group)	3		Except mushrooms and sweet corn
	Ginseng	0.1		
	Ginseng, dried including red ginseng	0.3		
	Ginseng, extracts	0.5		
	Grapes	2		
	Guava	0.2		
	Herbs	70		
	Hops, dry	30		

	Legume vegetables (group)	3		
	Lettuce, head	3		
	Lettuce, leaf	3		
	Maize	0.03	*	
	Maize oil, edible	0.1		
	Mango	0.7		
	Meat (from mammals other than marine mammals)	0.05		fat
	Milk fats	0.03		
	Milks	0.01		
	Oats	1.5		
	Papaya	0.3		
	Peanut	0.2		
	Peppers chili, dried	30		
	Pistachio nuts	1		
	Pitaya	0.3		
	Plantain	2		
	Potato	7		Po
	Poultry meat	0.01	(*)	
	Poultry, edible offal of	0.01	(*)	
	Pulses (group)	0.07		(except soya beans)
	Rape seed	0.5		
	Rice	5		
	Root and tuber vegetables (group)	1		(except potato)
	Rye	0.2		
	Sorghum grain	10		
	Soya bean (dry)	0.5		
	Stone fruits (group)	2		
	Straw and hay of cereal grains except pseudocereals	15		dry wt (except maize and sorghum)
	Strawberry	10		
	Sugar cane	0.05		
	Sunflower seed	0.5		
	Tree nuts (group)	0.01		
	Triticale	0.2		
	Wheat	0.2		
	Witloof chicory (sprouts)	0.3		
	Cumin	0.03	*	
Benalaxyl	Grapes	0.3		
	Lettuce, head	1		
	Melons, except watermelon	0.3		
	Onion, bulb	0.02	(*)	

	Potato	0.02	(*)	
	Tomato	0.2		
	Watermelon	0.1		
Bensulfuron Methyl	Rice	0.01		
Bentazone	Barley, hay and/or straw	0.3		
	Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	0.01	(*)	(green pods and immature seeds)
	Beans, shelled	0.01	(*)	(succulent=immature seeds)
	Cereal grains	0.01	(*)	
	Dry beans (subgroup)	0.5		
	Dry peas (subgroup)	0.5		
	Edible offal (mammalian)	0.04		
	Eggs	0.01	(*)	
	Herbs	0.1		
	Linseed	0.02	(*)	
	Mammalian fats (except milk fats)	0.01	(*)	
	Meat (from mammals other than marine mammals)	0.01		
	Milks	0.01	(*)	
	Onion, bulb	0.04		
	Peanut	0.05	(*)	
	Peas (pods and succulent=immature seeds)	1.5		
	Potato	0.1		
	Poultry meat	0.03		fat
	Poultry, edible offal of	0.07		
	Spring onion	0.08		
	Sweet corn (corn-on-the- cob)	0.01		
	Triticale, hay and/or straw	0.3		
	Wheat, hay and/or straw	0.3		
	Rice	0.05	*	
Cyfluthrin/beta-cyfluthrin	Soybean	0.03		
	Soybean oil	0.01*		
	Apple	0.1		
	Cabbages, head	0.08		
	Cauliflower	2		
	Citrus fruits (group)	0.3		
	Citrus pulp, dried	2		
	Cotton seed	0.7		
	Cotton seed oil, crude	1		
	Edible offal (mammalian)	0.02		
	Egg plant	0.2		

	Eggs	0.01	(*)	
	Meat (from mammals other than marine mammals)	0.2		(fat)
	Milks	0.01		
	Pear	0.1		
	Peppers (subgroup)	0.2		
	Peppers chili, dried	1		
	Potato	0.01	(*)	
	Poultry meat	0.01	(*)	(fat)
	Poultry, edible offal of	0.01	(*)	
	Rape seed	0.07		
	Soya bean (dry)	0.03		
	Spices, fruits and berries	0.03		
	Spices, roots and rhizomes	0.05		
	Tomato	0.2		
Bifenazate	Almond hulls	10		
	Beans (dry)	0.3		
	Blackberries	7		
	Cotton seed	0.3		
	Dewberries (including boysenberry and loganberry)	7		
	Edible offal (mammalian)	0.01	(*)	
	Eggs	0.01	(*)	
	Fruiting vegetables, cucurbits (group)	0.5		
	Grape, dried (= Currants, Raisins and Sultanas)	2		
	Grapes	0.7		
	Hops, dry	20		
	Legume vegetables (group)	7		
	Meat (from mammals other than marine mammals)	0.05		fat
	Milk fats	0.05		
	Milks	0.01	(*)	
	Mints	40		
	Peppers chili	3		
	Peppers, sweet (including pimento or pimienta)	2		
	Pome fruits (group)	0.7		
	Poultry meat	0.01	(*)	fat
	Poultry, edible offal of	0.01	(*)	
	Raspberries, red, black	7		
	Stone fruits (group)	2		
	Strawberry	2		

Bifenthrin	Tomato	0.5		
	Tree nuts (group)	0.2		
	Banana	0.1		
	Blackberries	1		
	Blueberries	3		
	Brassica vegetables (except Brassica leafy vegetables) (group)	0.4		
	Citrus fruits (group)	0.05		
	Cotton seed	0.5		
	Dewberries (including boysenberry and loganberry)	1		
	Edible offal (mammalian)	0.2		
	Egg plant	0.3		
	Grapes	0.3		
	Hops, dry	20		
	Maize	0.05	(*)	
	Meat (from mammals other than marine mammals)	3		fat
	Milk fats	3		
	Milks	0.2		
	Mustard greens	4		
	Peas (pods and succulent=immature seeds)	0.9		
	Peas, shelled (succulent seeds)	0.05	(*)	
	Peppers (subgroup)	0.5		
	Peppers chili, dried	5		
	Pulses (group)	0.3		
	Radish leaves (including radish tops)	4		
	Rape seed	0.05		
	Rape seed oil, edible	0.1		
	Raspberries, red, black	1		
	Root and tuber vegetables (group)	0.05		
	Spices, fruits and berries	0.03		
	Spices, roots and rhizomes	0.05		
	Straw and hay of cereal grains except pseudocereals	1		dw
	Tea, green, black (black, fermented and dried)	30		
	Tomato	0.3		
	Tree nuts (group)	0.05		
	Wheat	0.5		Po
	Wheat bran, unprocessed	2		PoP

	Wheat germ	1		Po
	Sugarcane	0.03		
	Rice	0.05		
	Apple	0.5		
Bispyribac Sodium	Rice	0.05		
Boscalid	Almond hulls	15		
	Banana	0.6		
	Barley	0.5		
	Barley, hay and/or straw	50		dry wt
	Berries and other small fruits	10		Except strawberry and grapes
	Brassica vegetables (except Brassica leafy vegetables) (group)	5		
	Bulb vegetables (group)	5		
	Cereal grains	0.1		Except barley, oats, rye and wheat
	Cherries (subgroup)	5		
	Citrus fruits (group)	2		
	Citrus oil, edible	50		
	Citrus pulp, dried	6		
	Coffee beans	0.05	(*)	
	Edible offal (mammalian)	0.2		
	Eggs	0.02		
	Fruiting vegetables, cucurbits (group)	3		
	Fruiting vegetables, other than cucurbits (group)	3		Except fungi, mushroom and sweet corn
	Grape, dried (= Currants, Raisins and Sultanas)	10		
	Grapes	5		
	Hops, dry	60		
	Kiwifruit	5		
	Leafy vegetables	40		
	Legume vegetables (group)	3		
	Mango	2		
	Meat (from mammals other than marine mammals)	0.7		fat
	Milk fats	2		
	Milks	0.1		
	Oats	0.5		
	Oilseed	1		

	Peaches (including apricots and nectarine) (subgroup)	4		
	Peppers chili, dried	10		
	Pistachio nuts	1		
	Plums (including fresh prunes) (subgroup)	1.5		
	Pome fruits (group)	2		
	Poultry fats	0.02		
	Poultry meat	0.02		
	Poultry, edible offal of	0.02		
	Prunes	5		
	Pulses (group)	3		
	Root and tuber vegetables (group)	2		
	Rye	0.5		
	Stalk and stem vegetables	30		
	Straw and hay of cereal grains except pseudocereals	5	dry wt	Except straw and fodder of barley, oats, rye and wheat
	Strawberry	3		
	Tea, green, black (black, fermented and dried)	40		
	Tree nuts (group)	0.05	(*)	Except pistachio
	Wheat	0.5		
	Wheat, hay and/or straw	50		dry wt
Buprofezin	Almond hulls	3		
	Apple	3		
	Avocado	0.1		
	Banana	0.3		
	Basil, leaves	1.5		
	Cherries (subgroup)	2		
	Citrus fruits (group)	1		
	Citrus oil, edible	6		
	Citrus pulp, dried	5		
	Coffee beans	0.4		
	Edible offal (mammalian)	0.05	(*)	
	Eggs	0.01	(*)	
	Fruiting vegetables, cucurbits (group)	0.7		
	Grape, dried (= Currants, Raisins and Sultanas)	2		
	Grapes	1		
	Mammalian fats (except milk fats)	0.01	(*)	
	Mango	0.1		

	Meat (from mammals other than marine mammals)	0.05	(*)	
	Milks	0.01	(*)	
	Nectarine	9		
	Olive oil, virgin	20		
	Peach	9		
	Pear	6		
	Peppers (subgroup)	2		
	Chilli	2		
	Dried Chilli	20		
	Peppers chili	10		
	Peppers chili, dried	10		
	Plums (including fresh prunes) (subgroup)	2		
	Poultry fats	0.01	(*)	
	Poultry meat	0.01	(*)	
	Poultry, edible offal of	0.01		
	Soya bean (dry)	0.01	(*)	
	Strawberry	3		
	Table olives	5		
	Tea, green	30		
	Tomato	1		
	Tree nuts (group)	0.05	(*)	
	Rice	0.05		
	Cotton seed oil	0.01		
Butachlor	Rice	0.05		
Cadusafos	Banana	0.01		
Carbaryl	Almond hulls	50		
	Asparagus	15		
	Beetroot	0.1		
	Carrot	0.5		
	Citrus fruits (group)	15		
	Cranberry	5		
	Egg plant	1		
	Kidney of cattle, goats, pigs and sheep	3		
	Liver of cattle, goats, pigs & sheep	1		
	Maize	0.5		
	Maize cob kernels	1		
	Maize oil, crude	0.1		
	Meat (from mammals other than marine mammals)	0.05		
	Milks	0.05		
	Olive oil, virgin	25		





	Chilli	5		
	Dried Chilli	50		
	Peppers, sweet (including pimento or pimiento)	5		
	Rice bran, unprocessed	170		
	Rice hulls	50		
	Rice, hay and/or straw	120		
	Rice	2.5		
	Sorghum forage (dry)	50		
	Sorghum grain	10		PoT
	Soya bean (dry)	0.2		
	Soya bean hulls	0.3		
	Soya bean oil, crude	0.2		
	Spices, fruits and berries	0.8		
	Spices, roots and rhizomes	0.1		
	Sunflower seed	0.2		
	Sunflower seed oil, crude	0.05		
	Sweet corn (corn-on-the-cob)	0.1		
	Sweet corn cannery waste	7.4		
	Sweet potato	0.02	(*)	
	Table olives	30		
	Tomato	5		
	Tomato juice	3		
	Tomato paste	10		
	Tree nuts (group)	1		
	Turnip, Garden	1		
	Wheat	2		
	Wheat bran, unprocessed	2		
	Wheat flour	0.2		
	Wheat germ	1		
	Wheat, hay and/or straw	30		
	Sesamum	0.05		
	Fish	0.2		
	Milled food grains	0.01		
Carbendazim	Apricot	2		
	Asparagus	0.2		
	Banana	0.2		
	Banana whole	1		
	Barley	0.5		
	Barley, hay and/or straw	2		
	Beans (dry)	0.5		
	Berries and other small fruits	1		Except grapes
	Brussels sprouts	0.5		
	Carrot	0.2		
	Cattle meat	0.05	(*)	

	Cherries (subgroup)	10		Based on thiophanatemethyl use
	Chicken fat	0.05	(*)	
	Coffee beans	0.1		
	Common bean (pods and/or immature seeds)	0.5		
	Cucumber	0.05	(*)	
	Edible offal (mammalian)	0.05	(*)	
	Cotton seed	0.1		
	Groundnut	0.1		
	Sugar beet	0.1		
	Eggs	0.1		(shell free basis)
	Garden pea, shelled (succulent seeds)	0.02		
	Gherkin	0.05	(*)	
	Grapes	3		
	Lettuce, head	5		
	Mango	5		Based on carbendazime use
	Milks	0.05	(*)	
	Nectarine	2		
	Oranges, sweet, sour (including Orange-like hybrids) (subgroup)	1		
	Peach	2		
	Peanut	0.1	(*)	
	Peppers chili	2		
	Peppers chili, dried	20		
	Pineapple	5		
	Plums (including fresh prunes) (subgroup)	0.5		
	Pome fruits (group)	3		
	Poultry meat	0.05	(*)	
	Rape seed	0.05	(*)	
	Rice, hay and/or straw	15		
	Rice, husked	2	(*)	
	Rye	0.1		
	Soya bean (dry)	0.5		
	Spices, fruits and berries	0.1		
	Spices, roots and rhizomes	0.1		
	Spices, seeds	5		
	Squash, summer	0.5		
	Sugar beet	0.1	(*)	
	Tomato	0.5		
	Tree nuts (group)	0.1	(*)	
	Wheat	0.05	(*)	

	Wheat, hay and/or straw	1		
	Tea	0.5		
	Meat and poultry	0.1		(Carcass fat basis)
	Other fruits	5		
	Vegetables	0.5		
	Milled food grains	0.1		
	other food grains	0.1		
Carbofuran (sum of carbofuran & 3-hydroxy carbofuran expressed as carbofuran)	Banana	0.01	(*)	
	Cattle fat	0.05	(*)	
	Citrus pulp, dried	2		Based on the use of carbosulfan
	Coffee beans	1		
	Cotton seed	0.1		
	Edible offal of cattle, goats, horses, pigs & sheep	0.05	(*)	
	Goat fat	0.05	(*)	
	Horse fat	0.05	(*)	
	Maize	0.05	(*)	Based on the use of carbosulfan
	Mandarin	0.5		Based on the use of carbosulfan
	Meat of cattle, goats, horses, pigs & sheep	0.05	(*)	
	Oranges, sweet, sour (including Orange-like hybrids) (subgroup)	0.5		
	Pig fat	0.05	(*)	
	Rape seed	0.05	(*)	
	Rice, hay and/or straw	1		
	Rice, husked	0.1		
	Sheep fat	0.05	(*)	
	Sorghum grain	0.1	(*)	
	Spices, roots and rhizomes	0.1		
	Sugar beet	0.2		Based on the use of carbosulfan
	Sugar cane	0.1	(*)	
	Sunflower seed	0.1	(*)	
Carbosulfan	Citrus pulp, dried	0.1		
	Cotton seed	0.05		
	Edible offal (mammalian)	0.05	(*)	
	Eggs	0.05	(*)	
	Maize	0.05	(*)	
	Mandarin	0.1		

	Meat (from mammals other than marine mammals)	0.05	(*) (fat)	
	Oranges, sweet, sour (including Orange-like hybrids) (subgroup)	0.1		
	Poultry meat	0.05	(*)	
	Poultry, edible offal of	0.05	(*)	
	Rice, hay and/or straw	0.05	(*)	
	Spices, fruits and berries	0.07		
	Spices, roots and rhizomes	0.1		
	Sugar beet	0.3		
	Chilli	2		
	Dried Chilli	20		
	Rice	0.2		
Carfentazone Ethyl	Wheat	0.01		
	Rice	0.1	*	
	Tea	0.02	*	
Cartap Hydrochloride	Rice	0.5		
Chlorantraniliprole	Artichoke, globe	2		
	Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	0.8		(green pods and immature seeds)
	Berries and other small fruits	1		
	Brassica vegetables (except Brassica leafy vegetables) (group)	2		
	Carrot	0.08		
	Celery	7		
	Cereal grains	0.02		(except rice)
	Citrus fruits (group)	0.7		
	Coffee beans	0.05		
	Cotton seed	0.3		
	Dry beans (subgroup)	0.3		(except soya beans)
	Dry peas (subgroup)	0.3		
	Edible offal (mammalian)	0.2		
	Eggs	0.2		
	Fruiting vegetables, cucurbits (group)	0.3		
	Fruiting vegetables, other than cucurbits (group)	0.6		
	Hops, dry	40		
	Leafy vegetables	20		(except radish leaves)
	Mammalian fats (except milk fats)	0.2		

	Meat (from mammals other than marine mammals)	0.2		fat
	Milk fats	0.2		
	Milks	0.05		
	Mints	15		
	Palm fruit (African oil palm)	0.8		
	Palm kernel oil, crude	2		
	Peanut	0.06		
	Peas (pods and succulent=immature seeds)	2		
	Peas, shelled (succulent seeds)	0.05		
	Maize	0.03	*	
	Chilli	0.6		
	Dried Chilli	6		
	Pome fruits (group)	0.4		
	Pomegranate	0.4		
	Poultry fats	0.08		
	Poultry meat	0.02		
	Poultry, edible offal of	0.07		
	Radish	0.5		
	Radish leaves (including radish tops)	40		
	Rape seed	2		
	Rice	0.4		
	Rice, polished	0.04		
	Root and tuber vegetables (group)	0.02		(except carrot and radish)
	Soya bean (dry)	0.05		
	Stone fruits (group)	1		
	Sugar cane	0.5		
	Sunflower seed	2		
	Sweet corn (corn-on-the-cob)	0.01	(*)	
	Tree nuts (group)	0.02		
	Groundnut	0.03	*	
	Groundnut oil	0.03	*	
Chlorfenapyr	Edible offal (mammalian)	0.05		
	Eggs	0.01		
	Garlic	0.01	(*)	
	Lemons and limes (including citron) (subgroup)	0.8		
	Mammalian fats (except milk fats)	0.6		

	Meat (from mammals other than marine mammals)	0.6		fat
	Melons, except watermelon	0.4		
	Milks	0.03		
	Onion, bulb	0.01	(*)	
	Oranges, sweet, sour (including Orange-like hybrids) (subgroup)	1.5		
	Papaya	0.3		
	Peppers (subgroup)	0.3		
	Peppers chili, dried	3		
	Potato	0.01	(*)	
	Poultry fats	0.02		
	Poultry meat	0.02		fat
	Poultry, edible offal of	0.01		
	Soya bean (dry)	0.08		
	Soya bean oil, crude	0.4		
	Tea, green, black (black, fermented and dried)	60		
	Tomato	0.4		
Chlorimuron ethyl	Rice	0.01		
	Soybean seed	0.01		
	Wheat	0.05		
Chlorothalonil	Asparagus	0.01	(*)	
	Banana	15		
	Brussels sprouts	6		
	Celery	20		
	Chard	50		
	Cherries (subgroup)	3		
	Cranberry	5		
	Cucumber	3		
	Currants, black, red, white	20		
	Edible offal (mammalian)	0.2		
	Flowerhead brassicas (includes Broccoli: Broccoli, Chinese and cauliflower)	5		
	Gherkin	3		
	Ginseng, dried including red ginseng	2		
	Gooseberry	20		
	Grapes	3		
	Horseradish	1		
	Leek	40		
	Mammalian fats (except milk fats)	0.07		

	Meat (from mammals other than marine mammals)	0.02		
	Melons, except watermelon	2		
	Milks	0.07		
	Onion, bulb	1.5		
	Onion, Chinese	10		
	Onion, Welsh	10		
	Papaya	20		
	Peaches (including apricots and nectarine) (subgroup)	1.5		
	Peanut	0.1		
	Peppers (subgroup)	7		
	Peppers chili, dried	70		
	Pistachio nuts	0.3		
	Poultry fats	0.01		
	Poultry meat	0.01		
	Poultry skin	0.01		
	Poultry, edible offal of	0.07		
	Pulses (group)	1		
	Rhubarb	7		
	Root and tuber vegetables (group)	0.3		(except horseradish)
	Shallot	1.5		
	Spring onion	10		
	Squash, summer	3		
	Strawberry	5		
	Tomato	5		
Chlorpropham	Cattle meat	0.1		fat
	Cattle, edible offal of	0.01	(*)	
	Milk fats	0.02		
	Milks	0.01	(*)	
	Potato	30		Po
Chlorpyrifos	Almonds	0.05		
	Banana	2		
	Broccoli	2		
	Cabbages, head	1		
	Carrot	0.1		
	Cattle kidney	0.01		
	Cattle liver	0.01		
	Cattle meat	1		fat
	Cauliflower	0.05		
	Chinese cabbage (type pe-tsai)	1		
	Citrus fruits (group)	1		
	Coffee beans	0.05		

Common bean (pods and/or immature seeds)	0.01		
Cotton seed	0.3		
Cotton seed oil, edible	0.05	(*)	
Cranberry	1		
Eggs	0.01	(*)	
Grape, dried (= Currants, Raisins and Sultanas)	0.1		
Grapes	0.5		
Maize	0.05		
Maize oil, edible	0.2		
Milk of cattle, goats & sheep	0.02		
Onion, bulb	0.2		
Peach	0.5		
Peas (pods and succulent=immature seeds)	0.01		
Pecan	0.05	(*)	
Peppers chili, dried	20		
Peppers, sweet (including pimento or pimienta)	2		
Pig meat	0.02		fat
Pig, edible offal of	0.01	(*)	
Plums (including fresh prunes) (subgroup)	0.5		
Pome fruits (group)	1		
Potato	2		
Poultry meat	0.01		fat
Poultry, edible offal of	0.01	(*)	
Rice	0.5		
Sheep meat	1		
Sheep, edible offal of	0.01		
Sorghum grain	0.5		
Soya bean (dry)	0.1		
Soya bean oil, refined	0.03		
Spices, fruits and berries	1		
Spices, roots and rhizomes	1		
Spices, seeds	5		
Strawberry	0.3		
Sugar beet	0.05		
Sweet corn (corn-on-the-cob)	0.01		
Tea, green, black (black, fermented and dried)	2		
Walnuts	0.05	(*)	
Wheat	0.5		
Wheat flour	0.1		



	Wheat, hay and/or straw	5		
	Food grains	0.05		
	Other vegetables	0.2		
	Carbonated Water	0.001		
Clodinafop-propargyl	Soybean	0.05	*	
	Wheat	0.1		
Clomazone	Rice	0.01		
	Soybean seed	0.01		
	Soybean seed oil	0.01		
Chlothianidin (Chlothianidin and its metabolites Thiazolymethylguanidine (TMG), Thiazolymethylurea (TZMU), Methylnitroguanidine (MNG) TMG)	Artichoke, globe	0.05		
	Avocado	0.03		
	Banana	0.02		
	Barley	0.04		
	Barley, hay and/or straw	0.2		dry wt
	Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	0.2		
	Berries and other small fruits	0.07		Except grapes
	Brassica vegetables (except Brassica leafy vegetables) (group)	0.2		
	Cacao beans	0.02	(*)	
	Celery	0.04		
	Citrus fruits (group)	0.07		
	Coffee beans	0.05		
	Edible offal (mammalian)	0.02	(*)	Except liver.
	Eggs	0.01	(*)	
	Fruiting vegetables, cucurbits (group)	0.02	(*)	
	Fruiting vegetables, other than cucurbits (group)	0.05		Except sweet corn
	Grape juice	0.2		
	Grape, dried (= Currants, Raisins and Sultanias)	1		
	Grapes	0.7		
	Hops, dry	0.07		
	Leafy vegetables	2		
	Legume vegetables (group)	0.01	(*)	
	Liver of cattle, goats, pigs & sheep	0.2		
	Maize	0.02		

	Mammalian fats (except milk fats)	0.02	(*)	
	Mango	0.04		
	Meat (from mammals other than marine mammals)	0.02	(*)	
	Milks	0.02		
	Mints	0.3		
	Oilseed	0.02	(*)	
	Papaya	0.01	(*)	
	Pecan	0.01	(*)	
	Peppers chili, dried	0.5		
	Pineapple	0.01	(*)	
	Pome fruits (group)	0.4		
	Popcorn	0.01	(*)	
	Poultry fats	0.01	(*)	
	Poultry meat	0.01	(*)	
	Poultry, edible offal of	0.1		
	Prunes	0.2		
	Pulses (group)	0.02		
	Rice	0.5		
	Root and tuber vegetables (group)	0.2		
	Sorghum grain	0.01	(*)	
	Stalk and stem vegetables	0.04		Except artichoke and celery
	Stone fruits (group)	0.2		
	Sugar cane	0.4		
	Sweet corn (corn-on-the-cob)	0.01	(*)	
	Tea, green, black (black, fermented and dried)	0.7		
	Wheat	0.02	(*)	
	Wheat, hay and/or straw	0.2		dry wt
Cyantraniliprole	Beans (dry)	0.3		
	Beans, shelled	0.3		
	Cotton seed	1.5		
	Cotton seed oil	1.5		
	Pomegranate seed	0.01		
	Brassica vegetables (except Brassica leafy vegetables) (group)	2		

Bush berries (subgroup)	4		
Celery	15		
Cherries (subgroup)	6		
Citrus fruits (group)	0.7		
Citrus oil, edible	4.5		
Coffee beans	0.05		
Common bean (pods and/or immature seeds)	1.5		
Cotton seed	1.5		
Cranberry	0.08		
Edible offal (mammalian)	1.5		
Eggs	0.15		
Fruiting vegetables, cucurbits (group)	0.3		
Fruiting vegetables, other than cucurbits (group)	0.5		(except mushrooms & sweet corn)
Garlic	0.05		
Leafy vegetables	20		(except Lettuce, Head)
Lettuce, head	5		
Maize	0.01	(*)	
Mammalian fats (except milk fats)	0.5		
Mango	0.7		
Meat (from mammals other than marine mammals)	0.2		
Milks	0.6		
Grapes	0.01		
Onion, bulb	0.05		
Onion, Welsh	8		
Peach	1.5		
Peas (pods and succulent=immature seeds)	2		
Peas, shelled (succulent seeds)	0.3		
Peppers chili, dried	5		
Plums (including fresh prunes) (subgroup)	0.5		
Pome fruits (group)	0.8		
Pomegranate	0.01	(*)	
Potato	0.05		
Poultry fats	0.04		
Poultry meat	0.02		
Poultry, edible offal of	0.15		
Prunes	0.8		

	Rape seed	0.8		
	Rice, hay and/or straw	1.7		dw
	Rice, husked	0.01	(*)	
	Rice, polished	0.01	(*)	
	Root and tuber vegetables (group)	0.05		(except potato)
	Shallot	0.05		
	Soya bean (dry)	0.4		
	Soya bean (immature seeds)	0.3		
	Spring onion	8		
	Strawberry	1.5		
	Sunflower seed	0.5		
	Tree nuts (group)	0.04		
	Wine-grapes	1		
Cyazofamid	Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	0.4		
	Beans, shelled	0.07		
	Brassica vegetables (except Brassica leafy vegetables) (group)	1.5		
	Bulb onions (subgroup)	1.5		
	Egg plant	0.2		
	Fruiting vegetables, cucurbits (group)	0.09		
	Grapes	1.5		
	Green onions (subgroup)	6		
	Hops, dry	15		
	Leafy vegetables	10		(except brassica leafy vegetables)
	Leaves of Brassicaceae	15		
	Peppers chili	0.8		
	Peppers, sweet (including pimento or pimienta)	0.4		
	Potato	0.01	(*)	
	Tomato	0.2		
Cyhalofop-butyl	Rice	0.5		
Cymoxanil	Tomato	0.01	*	
	Potato	0.01		
	Grapes	0.1		
	Citrus	0.05	*	
	Gherkin	0.05	*	
	Cucumber	0.1		
Cypermethrins (including alpha- and zeta-cypermethrin)	Artichoke, globe	0.1		

Asparagus	0.4		
Barley	2		Po
Brassica vegetables (except Brassica leafy vegetables) (group)	1		
Carambola	0.2		
Cardamom	3		
Cereal grains	0.3		Except rice barley, oats, rye and wheat.
Citrus fruits (group)	0.3		(excluding shaddocks or pomelos)
Coffee beans	0.05	(*)	
Durian	1		
Edible offal (mammalian)	0.05	(*)	The MRL accommodates external animal treatment.
Egg plant	0.03		
Eggs	0.01	(*)	
Fruiting vegetables, cucurbits (group)	0.07		
Ginseng	0.03	(*)	
Ginseng, dried including red ginseng	0.15		
Ginseng, extracts	0.06		
Grape, dried (= Currants, Raisins and Sultanas)	0.5		
Grapes	0.2		
Leafy vegetables	0.7		
Leek	0.05		
Legume vegetables (group)	0.7		



	Litchi	2		
	Longan	1		
	Mango	0.7		
	Meat (from mammals other than marine mammals)	2	fat	The MRL accommodates external animal treatment.
	Milk fats	0.5		
	Milks	0.05		The MRL accommodates external animal treatment.
	Oats	2		Po
	Oilseed	0.1		
	Okra	0.5		
	Olive oil, refined	0.5		
	Olive oil, virgin	0.5		
	Onion, bulb	0.01	(*)	
	Papaya	0.5		
	Peppers chili	2		
	Peppers chili, dried	10		
	Peppers, sweet (including pimento or pimiento)	0.1		
	Pome fruits (group)	0.7		
	Poultry fats	0.1		
	Poultry meat	0.1		fat
	Poultry, edible offal of	0.05	(*)	
	Pulses (group)	0.05	(*)	



	Pummelo and grapefruits (including Shaddock-like hybrids, among others Grapefruit) (subgroup)	0.5		
	Rice	2		
	Root and tuber vegetables (group)	0.01	(*)	Except sugar beet
	Rye	2		Po
	Spices, fruits and berries	0.5		
	Spices, roots and rhizomes	0.2		
	Stone fruits (group)	2		
	Straw and hay of cereal grains except pseudocereals	10		
	Strawberry	0.07		
	Sugar beet	0.1		
	Sugar cane	0.2		
	Sweet corn (corn-on-the-cob)	0.05	(*)	
	Table olives	0.05	(*)	
	Tea, green, black (black, fermented and dried)	15		
	Tomato	0.2		
	Tree nuts (group)	0.05	(*)	
	Wheat	2		Po
	Wheat bran, unprocessed	5		PoP
Cyproconazole	Beans (dry)	0.02	(*)	
	Cereal grains	0.08		Except maize, rice and sorghum
	Coffee beans	0.07		
	Coffee beans, roasted	0.1		
	Edible offal (mammalian)	0.5		
	Eggs	0.01	(*)	
	Maize	0.01	(*)	

	Meat (from mammals other than marine mammals)	0.02		fat
	Milks	0.01		
	Peas (dry)	0.02	(*)	
	Peas, shelled (succulent seeds)	0.01		
	Poultry meat	0.01	(*)	
	Poultry, edible offal of	0.01	(*)	
	Rape seed	0.4		
	Soya bean (dry)	0.07		
	Soya bean oil, refined	0.1		
	Straw and hay of cereal grains except pseudocereals	5		Except maize, rice and sorghum
	Sugar beet	0.05		
Cyromazine	Artichoke, globe	3		
	Beans (dry)	3		
	Broccoli	1		
	Celery	4		
	Chick-pea (dry)	3		
	Cucumber	2		
	Edible offal (mammalian)	0.3		
	Eggs	0.3		
	Fruiting vegetables, other than cucurbits (group)	1		Except mushrooms and sweet corn (corn-on-the-cob)
	Lentil (dry)	3		
	Lettuce, head	4		
	Lettuce, leaf	4		
	Lima bean (young pods and/or immature beans)	1		
	Lupin (dry)	3		
	Mango	0.5		
	Meat (from mammals other than marine mammals)	0.3		
	Melons, except watermelon	0.5		
	Milks	0.01		
	Mushrooms	7		
	Mustard greens	10		
	Onion, bulb	0.1		
	Peppers chili, dried	10		
	Poultry meat	0.1		
	Poultry, edible offal of	0.2		
	Spring onion	3		







Deltamethrin	Squash, summer	2		
	Apple	0.2		
	Carrot	0.02		
	Cereal grains	2	Po	
	Citrus fruits (group)	0.02		
	Eggs	0.02	(*)	
	Flowerhead brassicas (includes Broccoli: Broccoli, Chinese and cauliflower)	0.1		
	Fruiting vegetables, cucurbits (group)	0.2		
	Grapes	0.2		
	Hazelnuts	0.02	(*)	
	Kidney of cattle, goats, pigs and sheep	0.03	(*)	
	Leafy vegetables	2		
	Leek	0.2		
	Legume vegetables (group)	0.2		
	Liver of cattle, goats, pigs & sheep	0.03	(*)	
	Meat (from mammals other than marine mammals)	0.5	(fat)	The MRL accommodates external animal treatment.
	Milks	0.05	F	
	Mushrooms	0.05		
	Nectarine	0.05		
	Onion, bulb	0.05		
	Peach	0.05		
	Plums (including fresh prunes) (subgroup)	0.05		
	Potato	0.01	(*)	
	Poultry meat	0.1	(fat)	
	Poultry, edible offal of	0.02	(*)	
	Pulses (group)	1	Po	
	Radish	0.01	(*)	
	Rape seed	0.2		
	Spices, fruits and berries	0.03		
	Spices, roots and rhizomes	0.5		
	Strawberry	0.2		
	Sunflower seed	0.05	(*)	
	Sweet corn (corn-on-the- cob)	0.02	(*)	
	Table olives	1		

	Tea, green, black (black, fermented and dried)	5		
	Tomato	0.3		
	Walnuts	0.02	(*)	
	Wheat bran, unprocessed	5	PoP	
	Wheat flour	0.3	PoP	
	Wheat wholemeal	2	PoP	
	Red gram	0.01		
	Mango	0.01		
	Tea	5		
	Chilli	0.05		
	Dried Chilli	0.5		
Diafenthuron	Cardamom	0.5		
	Brinjal	1		
	Chilli	0.05		
	Dried Chilli	0.5		
	Cottonseed Oil	1		
	Cabbage	1		
	Citrus	0.2		
Diazinon	Almond hulls	5		
	Almonds	0.05		
	Blackberries	0.1		
	Boysenberry	0.1		
	Broccoli	0.5		
	Cabbages, head	0.5		
	Cantaloupe	0.2		
	Carrot	0.5		
	Cherries (subgroup)	1		
	Chicken eggs	0.02	(*)	
	Chicken meat	0.02	(*)	
	Chicken, edible offal of	0.02	(*)	
	Chinese cabbage (type pe-tsai)	0.05		
	Common bean (pods and/or immature seeds)	0.2		
	Cranberry	0.2		
	Cucumber	0.1		
	Currants, black, red, white	0.2		
	Garden pea, shelled (succulent seeds)	0.2		
	Goat meat	2	(fat)V	The MRL accommodates external animal treatment.
	Hops, dry	0.5		

Kale (Collards, curly kale, Scotch kale, thousand-headed kale; not including Marrow-stem kele)	0.05		
Kidney of cattle, goats, pigs and sheep	0.03	V	The MRL accommodates external animal treatment.
Kiwifruit	0.2		
Kohlrabi	0.2		
Lettuce, head	0.5		
Lettuce, leaf	0.5		
Liver of cattle, goats, pigs & sheep	0.03	V	The MRL accommodates external animal treatment.
Maize	0.02	(*)	
Meat of cattle, pigs & sheep	2	(fat)V	The MRL accommodates external animal treatment.
Milks	0.02	F	The MRL accommodates external animal treatment.
Onion, bulb	0.05		
Peach	0.2		
Peppers chili, dried	0.5		
Peppers, sweet (including pimento or pimienta)	0.05		
Pineapple	0.1		
Plums (including fresh prunes) (subgroup)	1		
Pome fruits (group)	0.3		
Potato	0.01	(*)	
Prunes	2		
Radish	0.1		
Raspberries, red, black	0.2		
Spices, fruits and berries	0.1	(*)	
Spices, roots and rhizomes	0.5		
Spices, seeds	5		
Spinach	0.5		
Spring onion	1		
Squash, summer	0.05		
Strawberry	0.1		
Sugar beet	0.1		

	Sweet corn (corn-on-the-cob)	0.02		
	Tomato	0.5		
	Walnuts	0.01	(*)	
Difenoconazole	Artichoke, globe	1.5		
	Asparagus	0.03		
	Avocado	0.6		
	Banana	0.1		
	Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	0.7		
	Blueberries	4		
	Brassica vegetables (except Brassica leafy vegetables) (group)	2		
	Carrot	0.2		
	Celeriac	0.5		
	Celery	3		
	Cherries (subgroup)	0.2		
	Citrus fruits (group)	0.6		
	Coffee beans	0.01	(*)	
	Cucumber	0.2		
	Dry beans (subgroup)	0.05		(except soya bean)
	Dry peas (subgroup)	0.15		
	Edible offal (mammalian)	1.5		
	Eggs	0.03		
	Fruiting vegetables, other than cucurbits (group)	0.6		(except peppers, chilli)
	Garlic	0.02	(*)	
	Gherkin	0.2		
	Ginseng	0.08		
	Ginseng, dried including red ginseng	0.8		
	Ginseng, extracts	0.6		
	Grape, dried (= Currants, Raisins and Sultanas)	6		
	Grapes	3		
	Leek	0.3		
	Lettuce, head	2		
	Lettuce, leaf	2		
	Mango	0.07		
	Meat (from mammals other than marine mammals)	0.2	(fat)	
	Melons, except watermelon	0.7		
	Milks	0.02		

	Nectarine	0.5		
	Onion, bulb	0.1		
	Papaya	0.2		
	Passion fruit	0.05		
	Peach	0.5		
	Peanut	0.01	(*)	
	Peas (pods and succulent=immature seeds)	0.7		
	Peppers chili	0.9		
	Peppers chili, dried	5		
	Pitaya	0.15		
	Plums (including fresh prunes) (subgroup)	0.2		
	Pome fruits (group)	4	Po	
	Potato	4	Po	
	Poultry meat	0.01	(*) (fat)	
	Poultry, edible offal of	0.01	(*)	
	Rape seed	0.15		
	Rice	8		
	Rice, hay and/or straw	17	(dw)	
	Rice, polished	0.07		
	Soya bean (dry)	0.1		
	Spring onion	9		
	Squash, summer	0.2		
	Strawberry	2		
	Pomegranate	0.8		
	Sugar beet	0.2		
	Sunflower seed	0.02		
	Sweet corn (corn on the cob) (kernels plus cob with husk removed)	0.01	(*)	
	Table olives	2		
	Tree nuts (group)	0.03		
	Watermelon	0.02		
	Wheat	0.02	(*)	
	Maize	0.01	*	
	Wheat, hay and/or straw	3		
Dimethoate	Artichoke, globe	0.05		
	Asparagus	0.05	(*)	
	Barley	2		
	Brussels sprouts	0.2		
	Cabbage, Savoy	0.05	(*)	
	Cattle, edible offal of	0.05	(*)	
	Cauliflower	0.2		
	Celery	0.5		
	Cherries (subgroup)	2		



	Citrus fruits (group)	5		(excluding kumquats)
	Eggs	0.05	(*)	
	Lettuce, head	0.3		
	Mammalian fats (except milk fats)	0.05	(*)	
	Mango	1	Po	
	Meat of cattle, goats, horses, pigs & sheep	0.05	(*)	
	Milk of cattle, goats & sheep	0.05	(*)	
	Pear	1		
	Peas (pods and succulent=immature seeds)	1		
	Chilli	0.5		
	Dried Chilli	5		
	Peppers, sweet (including pimento or pimiento)	0.5		
	Potato	0.05		
	Poultry fats	0.05	(*)	
	Poultry meat	0.05	(*)	
	Poultry, edible offal of	0.05	(*)	
	Sheep, edible offal of	0.05	(*)	
	Spices, fruits and berries	0.5		
	Spices, roots and rhizomes	0.1	(*)	
	Spices, seeds	5		
	Sugar beet	0.05		
	Table olives	0.5		
	Turnip greens	1		
	Turnip, Garden	0.1		
	Wheat	0.05		
	Wheat, hay and/or straw	1		
	Mustard	0.01		
	Fruits	2		
	Vegetables	2		
Dimethomorph	Artichoke, globe	2		
	Beans, shelled	0.7		
	Broccoli	4		
	Cabbages, head	6		
	Celery	15		
	Corn salad	10		
	Edible offal (mammalian)	0.01	(*)	
	Eggs	0.01	(*)	
	Fruiting vegetables, cucurbits (group)	0.5		



	Fruiting vegetables, other than cucurbits (group)	1.5		
	Garlic	0.6		
	Grape, dried (= Currants, Raisins and Sultanas)	5		
	Grapes	3		
	Hops, dry	80		
	Kohlrabi	0.02		
	Leek	0.8		
	Lettuce, head	10		
	Lettuce, leaf	9		
	Meat (from mammals other than marine mammals)	0.01	(*)	
	Milks	0.01	(*)	
	Onion, bulb	0.6		
	Onion, Welsh	9		
	Peas, shelled (succulent seeds)	0.15		
	Peppers chili, dried	5		
	Pineapple	0.01	(*)	
	Potato	0.05		
	Poultry meat	0.01	(*)	
	Poultry, edible offal of	0.01	(*)	
	Shallot	0.6		
	Spinach	30		
	Spring onion	9		
	Strawberry	0.5		
	Taro leaves	10		
Dinotefuran	Brassica vegetables (except Brassica leafy vegetables) (group)	2		
	Celery	0.6		
	Cotton seed	0.2		
	Cranberry	0.15		
	Edible offal (mammalian)	0.1		
	Eggs	0.02	(*)	
	Fruiting vegetables, cucurbits (group)	0.5		
	Fruiting vegetables, other than cucurbits (group)	0.5		(except sweet corn and mushrooms)
	Grape, dried (= Currants, Raisins and Sultanas)	3		
	Grapes	0.9		
	Leafy vegetables	6		(except watercress)

	Meat (from mammals other than marine mammals)	0.1		
	Milks	0.1		
	Nectarine	0.8		
	Onion, bulb	0.1		
	Peach	0.8		
	Peppers chili, dried	5		
	Poultry meat	0.02	(*)	
	Poultry, edible offal of	0.02	(*)	
	Rice	8		
	Rice, hay and/or straw	6		
	Rice, polished	0.3		
	Spring onion	4		
	Watercress	7		
	Cottonseed Oil	0.05	*	
Diuron	Sugarcane	0.02		
	Cottonseed	1		
	Banana	0.1		
	Maize	0.5		
	Citrus (Sweet Orange)	1		
	Grapes	1		
Edifenphos	Rice	0.02		
	Rice bran	1		
	Eggs	0.01		(shell free basis)
	Meat and poultry	0.02		(carcass fat basis)
	Milks	0.01		(F)
Emamectin benzoate	Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	0.01		
	Cos lettuce	0.7		
	Edible offal (mammalian)	0.08		
	Fruiting vegetables, cucurbits (group)	0.007		
	Fruiting vegetables, other than cucurbits (group)	0.02		except sweet corn and mushrooms
	Grapes	0.03		
	Lettuce, head	1		
	Lettuce, leaf	0.7		
	Mammalian fats (except milk fats)	0.02		
	Meat (from mammals other than marine mammals)	0.004		
	Mustard greens	0.2		
	Nectarine	0.03		
	Peach	0.03		

	Peppers chili, dried	0.2		
	Pome fruits (group)	0.02		
	Rape seed	0.005	(*)	
	Tree nuts (group)	0.001	(*)	
	Cottonseed	0.02		
	Cottonseed oil	0.02		
	Okra	0.05		
	Groundnut oil	0.05		
	Milks	0.01	*	
	Tea	0.01	*	
Epoxyconazole	Ground nut oil	0.05	*	
	Groundnut cake	0.05	*	
	Maize	0.01	*	
	cumin	0.01	*	
	coffee	0.05	*	
	wheat	0.01	*	
	soybean	0.05	*	
	soybean oil	0.05	*	
	Rice	0.05	*	
Esfenvalerate	Commodity	MRL		Note
	Cotton seed	0.05		
	Eggs	0.01	(*)	
	Poultry meat	0.01	(*) (fat)	
	Poultry, edible offal of	0.01	(*)	
	Rape seed	0.01	(*)	
	Tomato	0.1		
	Wheat	0.05		
	Wheat, hay and/or straw	2		
Ethion(Residues to be determined as ethion and its oxygen analogue and expressed as ethion)	Spices, fruits and berries	5		
	Spices, roots and rhizomes	0.3		
	Spices, seeds	3		
	Gram	0.01		
	Pigeon Pea	0.01		
	Soybean Seed	0.01		

	Tea	5		
	Cucumber	0.5		
	Squash	0.5		
	Other Vegetables	1		
	Cotton seed	0.5		
	Milks	0.5		(F)
	Meat and Poultry	0.2		(Carcass fat basis)
	Eggs	0.2		(shell free basis)
	Dry fruits	0.1		(shell free basis)
	Food grains	0.03		
	Milled food grains	0.01		
	Peaches	1		
	Other fruits	2		
Ethiprole	Commodity	MRL		Note
	Coffee beans	0.07		
	Coffee beans, roasted	0.2		
	Edible offal (mammalian)	0.1		
	Eggs	0.05		
	Mammalian fats (except milk fats)	0.15		
	Meat (from mammals other than marine mammals)	0.15	(fat)	

	Milk fats	0.5		
	Milks	0.01		
	Poultry fats	0.05		
	Poultry meat	0.05	(fat)	
	Poultry, edible offal of	0.05		
	Rice	3		
	Rice, husked	1.5		
	Rice, polished	0.4		
Ethoprophos	Banana	0.02		
	Cucumber	0.01		
	Edible offal (mammalian)	0.01	(*)	
	Meat (from mammals other than marine mammals)	0.01	(*)	
	Melons, except watermelon	0.02		
	Milks	0.01	(*)	
	Peppers chili, dried	0.2		
	Peppers, sweet (including pimento or pimienta)	0.05		
	Potato	0.05		
	Strawberry	0.02	(*)	
	Sugar cane	0.02		
	Sweet potato	0.05		
	Tomato	0.01	(*)	
	Turnip, Garden	0.02	(*)	
Ethoxysulfuron	Rice	0.01		
Etoxazole	Almond hulls	3		
	Citrus fruits (group)	0.1		
	Cucumber	0.02		
	Edible offal (mammalian)	0.01	(*)	
	Grapes	0.5		
	Hops, dry	15		
	Meat (from mammals other than marine mammals)	0.01	(*) (fat)	
	Milks	0.01	(*)	
	Mints	15		
	Pome fruits (group)	0.07		
	Tea, green, black (black, fermented and dried)	15		
	Tree nuts (group)	0.01	(*)	
	Brinjal	0.2		
Fenamidone	Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	0.8		
	Beans, shelled	0.15		

	Cabbages, head	0.9		
	Carrot	0.2		
	Celery	40		
	Cotton seed	0.02	(*)	
	Edible offal (mammalian)	0.01	(*)	
	Eggs	0.01	(*)	
	Flowerhead brassicas (includes Broccoli: Broccoli, Chinese and cauliflower)	4		
	Fruiting vegetables, cucurbits (group)	0.2		
	Fruiting vegetables, other than cucurbits (group)	1.5		(except chilli pepper, fungi, sweet corn)
	Garlic	0.15		
	Grapes	0.6		
	Leek	0.3		
	Lettuce, head	20		
	Lettuce, leaf	0.9		
	Meat (from mammals other than marine mammals)	0.01	(*)	fat
	Milk fats	0.02		
	Milks	0.01	(*)	
	Onion, bulb	0.15	(*)	
	Onion, Welsh	3	(*)	fat
	Peppers chili	4	(*)	
	Peppers chili, dried	30		
	Potato	0.02		
	Poultry fats	0.01		
	Poultry meat	0.01		
	Poultry, edible offal of	0.01		
	Shallot	0.15		
	Spring onion	3		
	Strawberry	0.04		
	Sunflower seed	0.02	(*)	
	Tomato ketchup	3		
	Tomato paste	4		
	Witloof chicory (sprouts)	0.01	(*)	
Fenazaquin	Almond hulls	4	(dw)	
	Cherries (subgroup)	2		
	Edible offal (mammalian)	0.02	(*)	
	Hops, dry	30		
	Mammalian fats (except milk fats)	0.02	(*)	

	Meat (from mammals other than marine mammals)	0.02	(*) (fat)	
	Milk fats	0.02	(*)	
	Milks	0.02	(*)	
	Tree nuts (group)	0.02		
	Apple	0.2		
	Chilli	0.5		
	Dried Chilli	5		
	Okra	0.01		
	Brinjal	0.01		
	Tomato	0.01		
	Tea	3		
Fenbutatin Oxide	Almonds	0.5		
	Banana	10		
	Cherries (subgroup)	10		
	Chicken meat	0.05	(*)	
	Chicken, edible offal of	0.05	(*)	
	Citrus fruits (group)	5		(including kumquats)
	Citrus pulp, dried	25		
	Cucumber	0.5		
	Edible offal (mammalian)	0.2		
	Eggs	0.05		
	Grape pomace, dry	100		
	Grapes	5		
	Meat (from mammals other than marine mammals)	0.05	(*)	
	Milks	0.05	(*)	
	Peach	7		
	Pecan	0.5		
	Plums (including fresh prunes) (subgroup)	3		
	Pome fruits (group)	5		
	Prunes	10		
	Raisins (seedless white grape var., partially dried)	20		
	Strawberry	10		
	Tomato	1		
	Walnuts	0.5		
Fenitrothion	Apple	0.5		
	Cereal grains	6	Po	
	Edible offal (mammalian)	0.05	(*)	
	Eggs	0.05	(*)	

	Meat (from mammals other than marine mammals)	0.05	(*)	
	Milks	0.01		
	Poultry meat	0.05	(*)	
	Rice bran, unprocessed	40	PoP	
	Soya bean (dry)	0.01		
	Spices, fruits and berries	1		
	Spices, roots and rhizomes	0.1	(*)	
	Spices, seeds	7		
	Wheat bran, unprocessed	25	PoP	
Fenpropathrin	Almond hulls	10		
	Citrus fruits (group)	2		
	Citrus oil, edible	100		
	Coffee beans	0.03		
	Edible offal (mammalian)	0.01		
	Eggs	0.01	(*)	
	Mammalian fats (except milk fats)	0.03		
	Peppers (subgroup)	1		
	Peppers chili, dried	10		
	Plums (including fresh prunes) (subgroup)	1		
	Poultry fats	0.01	(*)	
	Poultry meat	0.01	(*) (fat)	
	Poultry, edible offal of	0.01	(*)	
	Prunes	3		
	Soya bean (dry)	0.01		
	Strawberry	2		
	Tea, green, black (black, fermented and dried)	3		
	Tomato	1		
	Tree nuts (group)	0.15		
	Brinjal	0.2		
	Okra	0.5		
	Rice	0.03	*	
	Cottonseed oil	3		
	Milks	0.1		
	Meats	0.02		
Fenpyroximate	Apple	0.2		
	Apples, dried	1		
	Avocado	0.2		
	Beans with pods (subgroup)	0.5		
	Cherries (subgroup)	2		
	Citrus fruits (group)	0.6		
	Citrus oil, edible	25		




	Coffee beans	0.07		
	Cucumber	0.3		
	Edible offal (mammalian)	0.5		
	Eggplants (subgroup)	0.3		
	Grape, dried (= Currants, Raisins and Sultanas)	0.2		
	Grapes	0.1		
	Hops, dry	15		
	Maize	0.01	(*)	
	Mammalian fats (except milk fats)	0.1		
	Meat (from mammals other than marine mammals)	0.1		fat
	Melons, except watermelon	0.2		
	Milks	0.01		
	Pear	0.2		
	Peppers (subgroup)	0.2		(except martynia, okra and roselle)
	Potato	0.05	(*)	
	Prunes	0.7		
	Raspberries, red, black	0.2		
	Squash, summer	0.06		
	Stone fruits (group)	0.4		(except cherries)
	Strawberry	0.3		
	Tea, green, black (black, fermented and dried)	8		
	Tomatoes (subgroup)	0.3		
	Tree nuts (group)	0.05	(*)	
	Chilli	1		
	Dried Chilli	10		
	Coconut Water	0.02		
Fenvalerate	Broccoli, Chinese	3		
	Edible offal (mammalian)	0.02		
	Mango	1.5		
	Meat (from mammals other than marine mammals)	1	(fat)	
	Milks	0.1	F	
	Spices, fruits and berries	0.03		
	Spices, roots and rhizomes	0.05		
	Cauliflower	2		
	Brinjal	2		
	Okra	2		

Fipronil	Cottonseed	0.2		
	Cottonseed Oil	0.1		
	Banana	0.005		
	Barley	0.002	(*)	
	Basil, leaves	1.5		
	Cabbages, head	0.02		
	Cattle kidney	0.02		
	Cattle liver	0.1		
	Cattle meat	0.5	(fat)	
	Cattle milk	0.02		
	Eggs	0.02		
	Flowerhead brassicas (includes Broccoli: Broccoli, Chinese and cauliflower)	0.02		
	Maize	0.01		
	Oats	0.002	(*)	
	Potato	0.02		
	Poultry meat	0.01	(*)	
	Poultry, edible offal of	0.02		
	Rice	0.01		
	Rice, hay and/or straw	0.2	dry wt	
	Rye	0.002	(*)	
	Sugar beet	0.2		
	Sunflower seed	0.002	(*)	
	Triticale	0.002	(*)	
	Cottonseed Oil	0.01		
	Rice	0.01		
	Chilli	0.01		
	Dried Chilli	0.1		
	Sugarcane	0.01		
	Cabbage	0.02		
	Grapes	0.01	*	
	wheat	0.01	*	
	Onion	0.04		
Flonicamid	Almond hulls	9		
	Almonds	0.01	(*)	
	Beans with pods (subgroup)	0.7		(except soya bean (succulent seeds in pods))
	Brassica vegetables (except Brassica leafy vegetables) (group)	2		
	Celery	1.5		
	Cherries (subgroup)	0.9		
	Citrus pulp, dried	3	(dw)	
	Cotton seed	0.6		

Rice	0.05	*	
Cottonseed Oil	0.02	*	
Dry beans (subgroup)	0.15		(except soya bean (dry))
Dry peas (subgroup)	1		
Edible offal (mammalian)	0.2		
Eggs	0.15		
Fruiting vegetables, cucurbits (group)	0.2		
Fruiting vegetables, other than cucurbits (group)	0.4		(except mushrooms and sweet corn)
Hops, dry	20		
Leaves of Brassicaceae	15		
Lemons and limes (including citron) (subgroup)	1.5		
Lettuce, head	1.5		
Lettuce, leaf	8		
Low growing berries (subgroup)	1.5		
Mammalian fats (except milk fats)	0.05		
Meat (from mammals other than marine mammals)	0.15		
Milks	0.15		
Mints	6		
Oranges, sweet, sour (including Orange-like hybrids) (subgroup)	0.4		
Peaches (including apricots and nectarine) (subgroup)	0.7		
Peas with pods (subgroup)	0.8		
Pecan	0.01	(*)	
Plums (including fresh prunes) (subgroup)	0.1		
Pome fruits (group)	0.8		
Potato	0.01		
Poultry fats	0.05		
Poultry meat	0.1		
Poultry, edible offal of	0.1		
Pummelo and grapefruits (including Shaddock-like hybrids, among others Grapefruit) (subgroup)	0.3		
Radish	0.4		

	Radish leaves (including radish tops)	20		
	Rape seed	0.5		
	Spinach	20		
	Succulent beans without pods (subgroup)	0.3		(except soya bean (succulent seeds))
	Succulent peas without pods (subgroup)	0.4		
	Tomato paste	7		
	Wheat	0.08		
	Wheat, hay and/or straw	0.3		
Fluazifop-p-butyl	Almonds	0.01	(*)	
	Banana	0.01	(*)	
	Beans (dry)	40		
	Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	6		
	Bush berries (subgroup)	0.3		
	Cabbages, head	3		
	Cane berries (subgroup)	0.08		
	Carrot	0.6		
	Celeriac	0.4		
	Citrus fruits (group)	0.01	(*)	
	Citrus pulp, dried	0.06	(*)	
	Coffee beans	0.01	(*)	
	Cotton seed	0.7		
	Edible offal (mammalian)	0.2		
	Egg plant	0.4		
	Eggs	0.03		
	Elderberries	0.3		
	Field pea (dry)	3		
	Garlic	0.3		
	Grapes	0.01	(*)	
	Guelder rose	0.3		
	Lettuce, leaf	0.01	(*)	
	Macadamia nuts	0.01	(*)	
	Mammalian fats (except milk fats)	0.09		
	Meat (from mammals other than marine mammals)	0.09	(fat)	
	Milks	0.2		
	Olives for oil production	0.01	(*)	
	Onion, bulb	0.3		
	Orange oil, edible	0.05	(*)	

	Peas (pods and succulent=immature seeds)	2		
	Peas, shelled (succulent seeds)	15		
	Pecan	0.01	(*)	
	Pome fruits (group)	0.01	(*)	
	Potato	0.6		
	Poultry fats	0.03		
	Poultry meat	0.03		
	Poultry, edible offal of	0.09		
	Shallot	0.3		
	Soya bean (dry)	15		
	Stone fruits (group)	0.01	(*)	
	Strawberry	3		
	Sugar beet	0.5		
	Sugar beet molasses	7		
	Sugar beet pulp, dry	20		
	Sugar cane	0.01	(*)	
	Sunflower seed	7		
	Swede	4		
	Table olives	0.01	(*)	
	Tomato	0.4		
	Turnip, Garden	4		
	Walnuts	0.01	(*)	
	Cotton seed Oil	0.01	*	
	Groundnut	0.01	*	
	Groundnut oil	0.01	*	
Flubendiamide	Almond hulls	10		
	Brassica vegetables (except Brassica leafy vegetables) (group)	4		
	Celery	5		
	Cotton seed	1.5		
	Edible offal (mammalian)	1		
	Fruiting vegetables, cucurbits (group)	0.2		
	Grapes	2		
	Legume vegetables (group)	2		
	Lettuce, head	5		
	Lettuce, leaf	7		
	Maize	0.02		
	Meat (from mammals other than marine mammals)	2	(fat)	
	Milk fats	5		
	Milks	0.1		
	Peppers (subgroup)	0.7		



	Peppers chili, dried	7		
	Pome fruits (group)	0.8		
	Pulses (group)	1		
	Stone fruits (group)	2		
	Sweet corn (corn-on-the-cob)	0.02		
	Tea, green, black (black, fermented and dried)	50		
	Tomato	2		
	Tree nuts (group)	0.1		
	Brinjal	0.1		
	Cottonseed Oil	1.5		
	Rice	0.1		
	Cabbage	4		
	Tomato	2		
	soyabean	0.07		
	soybean oil	0.07		
	soybean cake	0.07		
Fludioxonil	Avocado	1.5		
	Beans (dry)	0.5		
	Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	0.6		(green pods and immature seeds)
	Beans, shelled	0.4		
	Blackberries	5		
	Blueberries	2		
	Broccoli	0.7		
	Bulb onions (subgroup)	0.5		
	Cabbages, head	0.7		
	Carrot	1		
	Celery	15		
	Cereal grains	0.05	(*)	
	Chick-pea (dry)	0.3		
	Citrus fruits (group)	10		Po
	Cotton seed	0.05	(*)	
	Currants, black, red, white	3		
	Dewberries (including boysenberry and loganberry)	5		
	Dried herbs	60		
	Edible offal (mammalian)	0.1		
	Egg plant	0.3		
	Eggs	0.02		
	Fruiting vegetables, cucurbits (group)	0.5		
	Ginseng	4		
	Grapes	2		

Green onions (subgroup)	0.8		
Guava	0.5		
Herbs	9		
Kiwifruit	15		Po
Leaves of Brassicaceae	15		(except radish leaves)
Lentil (dry)	0.3		
Lettuce, head	10		
Lettuce, leaf	40		
Mammalian fats (except milk fats)	0.02		
Mango	2		
Meat (from mammals other than marine mammals)	0.02		fat
Milks	0.04		
Peas (dry)	0.07		
Peas (pods and succulent=immature seeds)	0.3		
Peas, shelled (succulent seeds)	0.03		
Peppers (subgroup)	1		
Peppers chili, dried	4		
Pineapple	5		Po
Pistachio nuts	0.2		
Pome fruits (group)	5		Po
Pomegranate	3		Po
Potato	5		Po
Poultry fats	0.01	(*)	
Poultry meat	0.01	(*)	
Poultry, edible offal of	0.1		
Radish	0.3		
Radish leaves (including radish tops)	20		
Rape seed	0.02	(*)	
Raspberries, red, black	5		
Snap bean (young pods)	0.6		
Soya bean (dry)	0.2		
Spinach	30		
Stone fruits (group)	5		Po
Straw and hay of cereal grains except pseudocereals	0.06	(*)	
Strawberry	3		
Sweet corn (corn-on-the-cob)	0.01	(*)	
Sweet potato	10		Po
Tomato	3		Po

	Watercress	10		
	Yams	10		Po
Flufenoxuron	Edible offal (mammalian)	0.05	(*)	
	Mammalian fats (except milk fats)	0.05	(*)	
	Meat (from mammals other than marine mammals)	0.05	(*)	
	Milks	0.01	(*)	
	Oranges, sweet, sour (including Orange-like hybrids) (subgroup)	0.4		
	Tea, green, black (black, fermented and dried)	20		
Fluopicolide	Brussels sprouts	0.2		
	Cabbages, head	7		
	Celery	20		
	Edible offal (mammalian)	0.01	(*)	
	Eggs	0.01	(*)	
	Flowerhead brassicas (includes Broccoli: Broccoli, Chinese and cauliflower)	2		
	Fruiting vegetables, cucurbits (group)	0.5		
	Fruiting vegetables, other than cucurbits (group)	1		Except mushrooms and sweet corn.
	Grape pomace, dry	7		
	Grape, dried (= Currants, Raisins and Sultanas)	10		
	Grapes	2		
	Leafy vegetables	30		
	Meat (from mammals other than marine mammals)	0.01	(*)	fat
	Milks	0.02		
	Onion, bulb	1		
	Onion, Welsh	10		
	Peppers chili, dried	7		
	Poultry meat	0.01	(*)	
	Poultry, edible offal of	0.01	(*)	
	Straw and hay of cereal grains except pseudocereals	0.2		
Fluopyram	Artichoke, globe	0.4		
	Asparagus	0.01	(*)	
	Banana	0.8		
	Barley	0.2		



Barley, hay and/or straw	2		
Basil, dry	400		
Basil, leaves	70		
Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	1		
Beans, shelled	0.2		
Broccoli	0.3		
Brussels sprouts	0.3		
Bush berries (subgroup)	7		
Cabbages, head	0.15		
Cane berries (subgroup)	5		
Carrot	0.4		
Cauliflower	0.09		
Cherries (subgroup)	2		
Cotton seed	0.8		
Cucumber	0.5		
Dill seed	70		
Dry beans (subgroup)	0.15		(except soy bean (dru))
Dry peas (subgroup)	0.7		
Edible offal (mammalian)	8		
Eggplants (subgroup)	0.5		
Eggs	2		
Garlic	0.07		
Grape, dried (= Currants, Raisins and Sultanas)	5		
Grapes	2		
Hops, dry	50		
Leek	0.15		
Lemons and limes (including citron) (subgroup)	1		
Lettuce, head	15		
Lettuce, leaf	15		
Maize cereals (subgroup)	0.02		
Mammalian fats (except milk fats)	1.5		
Mandarins (including mandarin-like hybrids) (subgroup)	0.6		
Mango	1		
Meat (from mammals other than marine mammals)	1.5		
Milks	0.8		
Oats	0.2		
Onion, bulb	0.07		

	Onion, Welsh	2		
	Oranges, sweet, sour (including Orange-like hybrids) (subgroup)	0.6		
	Peaches (including apricots and nectarine) (subgroup)	1		
	Peanut	0.2		
	Peas, shelled (succulent seeds)	0.2		
	Peppers (subgroup)	3		(except martynia, okra, roselle)
	Peppers chili, dried	30		
	Plums (including fresh prunes) (subgroup)	0.5		
	Pome fruits (group)	0.5		
	Potato	0.15		
	Poultry fats	1		
	Poultry meat	1.5		
	Poultry, edible offal of	5		
	Pummelo and grapefruits (including Shaddock-like hybrids, among others Grapefruit) (subgroup)	0.4		
	Rape seed	1		
	Rice	4		
	Rice, hay and/or straw	17		
	Rice, husked	1.5		
	Rice, polished	0.5		
	Rye	0.9		
	Soya bean (dry)	0.3		
	Spring onion	15		
	Strawberry	0.4		
	Sugar beet	0.04		
	Sunflower seed	0.7		
	Sweet corn (corn on the cob) (kernels plus cob with husk removed)	0.01	(*)	
	Tomatoes (subgroup)	0.5		
	Tree nuts (group)	0.04		
	Triticale	0.9		
	Triticale, hay and/or straw	23		
	Wheat	0.9		
	Wheat, hay and/or straw	23		
	Witloof chicory (sprouts)	0.15		
Flupyradiflurone and its metabolites Difluroacetic Acid and Difluroethylamino- furanone	Alfalfa hay	30	(dw)	

Apples, dried	2		
Avocado	0.6		
Bean hay	30		
Beans (dry)	0.4		
Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	1.5		(green pods and immature seeds)
Beans, shelled	0.2		(succulent = immature seeds)
Bulb vegetables, except fennel, bulb	0.01	(*)	
Bush berries (subgroup)	4		
Cabbages, head	1.5		
Cacao beans	0.01	(*)	
Cane berries (subgroup)	6		
Cauliflower	6		
Cereal grains	3		(except maize and rice)
Cherries (subgroup)	2		
Coffee beans	0.9		
Cotton seed	0.8		
Cucumber	0.4		
Edible offal (mammalian)	4		
Eggs	0.7		
Grape, dried (= Currants, Raisins and Sultanas)	8		
Grapes	3		



	Hops, dry	10		
	Lemons and limes (including citron) (subgroup)	1.5		
	Lettuce, head	4		
	Maize	0.01		
	Maize bran, unprocessed	0.05		
	Mammalian fats (except milk fats)	1		
	Mandarins (including mandarin-like hybrids) (subgroup)	1.5		
	Meat (from mammals other than marine mammals)	1.5		
	Melons, except watermelon	0.4		
	Milks	0.7		
	Oranges, sweet, sour (including Orange-like hybrids) (subgroup)	4		
	Pea hay	50	(dw)	
	Peaches (including apricots and nectarine) (subgroup)	1.5		
	Peanut	0.04		
	Peanut hay	30	(dw)	
	Peas (dry)	3		



	Peas (pods and succulent=immature seeds)	3		
	Peas, shelled (succulent seeds)	3		
	Pecan	0.01		
	Peppers (subgroup)	0.9		
	Peppers chili, dried	9		
	Plums (including fresh prunes) (subgroup)	0.4		
	Pome fruits (group)	0.9		
	Potato	0.05		
	Poultry fats	0.3		
	Poultry meat	0.8		
	Poultry, edible offal of	1		
	Prunes	3		
	Pummelo and grapefruits (including Shaddock-like hybrids, among others Grapefruit) (subgroup)	0.7		
	Root and tuber vegetables (group)	0.7		(except potato)
	Soya bean (dry)	1.5		



	Soya bean hay	40	(dw)	
	Squash, summer	0.2		
	Straw and hay of cereal grains except pseudocereals	40	(dw)	
	Strawberry	1.5		
	Sweet corn (corn-on-the-cob)	0.05		
	Sweet potato	0.05		
	Tomato	1		
	Wheat bran, unprocessed	8		
	Wheat germ	5		
	Wheat wholemeal	5		
	Okra	0.8		
Flusilazole	Apricot	0.2		
	Banana	0.03		
	Cereal grains	0.2		Except rice
	Edible offal (mammalian)	2		
	Eggs	0.1		
	Grape pomace, dry	2		
	Grape, dried (= Currants, Raisins and Sultanias)	0.3		
	Grapes	0.2		
	Meat (from mammals other than marine mammals)	1		fat
	Milks	0.05		
	Nectarine	0.2		



	Peach	0.2		
	Pome fruits (group)	0.3		
	Poultry meat	0.2		
	Poultry, edible offal of	0.2		
	Rape seed	0.1		
	Soya bean (dry)	0.05		
	Soya bean hulls	0.05		
	Soya bean oil, refined	0.1		
	Straw and hay of cereal grains except pseudocereals	5		Except rice
	Sugar beet	0.05		
	Sunflower seed	0.1		
	Sweet corn (corn-on-the-cob)	0.01	(*)	
	Rice	0.01		
	Chilli	0.01		
	Dried Chilli	0.1		
	Apple	0.05		
	Groundnut	0.05	*	
Fosetyl Al	Avocado	20		
	Blackberries	70		(based on Fosetyl-Al use)
	Coffee beans	30		(based on Fosetyl-Al use)
	Cucumber	60		
	Edible offal (mammalian)	0.5		
	Eggs	0.05	(*)	
	Flowerhead brassicas (subgroup)	0.2	(*)	(based on Fosetyl use)
	Grapes	60		
	Head brassicas (subgroup)	0.2	(*)	(based on Fosetyl use)
	Hops, dry	1500		
	Kale (Collards, curly kale, Scotch kale, thousand-headed kale; not including Marrow-stem kale)	0.2	(*)	(based on Fosetyl use)
	Kiwifruit	150		(based on Fosetyl-Al use)
	Lettuce, head	200		
	Lettuce, leaf	40		
	Mammalian fats (except milk fats)	0.3		
	Mandarins (including mandarin-like hybrids) (subgroup)	50		

	Meat (from mammals other than marine mammals)	0.15		
	Melons, except watermelon	60		
	Milks	0.1		
	Oranges, sweet, sour (including Orange-like hybrids) (subgroup)	20		
	Peppers, sweet (including pimento or pimienta)	7		
	Pineapple	15		(based on Fosetyl-Al use)
	Pome fruits (group)	50		
	Poultry fats	0.05	(*)	
	Poultry meat	0.05	(*)	
	Poultry, edible offal of	0.05	(*)	
	Spinach	20		
	Squash, summer	70		
	Strawberry	70		
	Tomato	8		
	Tree nuts (group)	400		
	Cardamom	0.2		
Glufosinate-Ammonium	Asparagus	0.4		
	Assorted tropical and sub-tropical fruits - edible peel	0.1		
	Assorted tropical and sub-tropical fruits - inedible peel	0.1		(except banana and kiwifruit)
	Banana	0.2		
	Blueberries	0.1		
	Carrot	0.05		
	Citrus fruits (group)	0.05		
	Coffee beans	0.1		
	Common bean (dry)	0.05		
	Common bean (pods and/or immature seeds)	0.05	(*)	
	Corn salad	0.05		
	Cotton seed	5		
	Currants, black, red, white	1		
	Edible offal (mammalian)	3		
	Eggs	0.05	(*)	
	Gooseberry	0.1		
	Grapes	0.15		
	Kiwifruit	0.6		
	Lettuce, head	0.4		



	Lettuce, leaf	0.4		
	Maize	0.1		
	Meat (from mammals other than marine mammals)	0.05		
	Milks	0.02	(*)	
	Onion, bulb	0.05		
	Pome fruits (group)	0.1		
	Potato	0.1		
	Poultry meat	0.05	(*)	
	Poultry, edible offal of	0.1	(*)	
	Prunes	0.3		
	Rape seed	1.5		
	Rape seed oil, crude	0.05	(*)	
	Raspberries, red, black	0.1		
	Rice	0.9		
	Rice, hay and/or straw	2		
	Soya bean (dry)	2		
	Stone fruits (group)	0.15		
	Strawberry	0.3		
	Sugar beet	1.5		
	Sugar beet molasses	8		
	Tree nuts (group)	0.1		
	Cottonseed Oil	0.05	*	
	Tea	0.01		
Glyphosate	Banana	0.05	(*)	
	Barley, hay and/or straw	400		
	Cereal grains	30		Except maize and rice
	Cotton seed	40		
	Dry beans (subgroup)	15		(except soya beans)
	Dry peas (subgroup)	10		
	Edible offal (mammalian)	5		Except pigs
	Eggs	0.05	(*)	
	Maize	5		
	Meat (from mammals other than marine mammals)	0.05	(*)	
	Milks	0.05	(*)	
	Pig, edible offal of	0.5		
	Poultry meat	0.05	(*)	
	Poultry, edible offal of	0.5		
	Rape seed	30		
	Soya bean (dry)	20		
	Sugar beet	15		
	Sugar cane	2		

	Sugar cane molasses	10		
	Sunflower seed	7		
	Sweet corn (corn-on-the-cob)	3		
	Wheat bran, unprocessed	20		
	Wheat, hay and/or straw	300		
	Tea	1		
	Rice	0.01		
Hexaconazole	Mango	0.02		
	Rice	0.02		
	Ground nut seed	0.02		
	Tea	0.02		
	Grapes	0.1		
	Chilli	0.5		
	Dried Chilli	5		
	Potato	0.02		
	Soybean	0.02		
	Apple	0.1		
	Black gram	0.01	*	
Hexythiazox	Citrus fruits (group)	0.5		
	Date	2		
	Edible offal (mammalian)	0.05		
	Egg plant	0.1		
	Eggs	0.05		
	Fruiting vegetables, cucurbits (group)	0.05		Except watermelon.
	Grape pomace, dry	15		
	Grape, dried (= Currants, Raisins and Sultanas)	1		
	Grapes	1		
	Hops, dry	3		
	Mammalian fats (except milk fats)	0.05		
	Meat (from mammals other than marine mammals)	0.05	(fat)	
	Milk fats	0.05		
	Milks	0.05		
	Pome fruits (group)	0.4		
	Poultry meat	0.05	(*) (fat)	
	Poultry, edible offal of	0.05		
	Prunes	1		
	Stone fruits (group)	0.3		
	Strawberry	6		
	Tea, green, black (black, fermented and dried)	15		
	Tomato	0.1		

	Tree nuts (group)	0.05	(*)	
	Chilli	0.01		
	Dried Chilli	0.1		
	Apple	0.3		
Imazethapyr	Edible offal (mammalian)	0.01	(*)	
	Eggs	0.01	(*)	
	Lentil (dry)	0.1	(*)	
	Maize	0.1	(*)	
	Mammalian fats (except milk fats)	0.01	(*)	
	Meat (from mammals other than marine mammals)	0.01	(*)	
	Milks	0.01	(*)	
	Peanut	0.1	(*)	
	Poultry fats	0.01	(*)	
	Poultry meat	0.01	(*)	
	Poultry, edible offal of	0.01	(*)	
	Rape seed	0.1	(*)	
	Rice	0.1	(*)	
	Rice, hay and/or straw	0.15	(*)(dw)	
	Soya bean (dry)	0.03	(*)	
	Soybean oil	0.1		
	Groundnut oil	0.1		
Imidacloprid	Almond hulls	5		
	Apple	0.5		
	Banana	0.05		
	Barley, hay and/or straw	1		
	Basil, leaves	20		
	Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	2		
	Berries and other small fruits	5		Except cranberries, grapes and strawberries.
	Broccoli	0.5		
	Brussels sprouts	0.5		
	Cabbages, head	0.5		
	Cauliflower	0.5		
	Celery	6		
	Cereal grains	0.05		
	Cherries (subgroup)	4		
	Citrus fruits (group)	1		
	Citrus pulp, dried	10		
	Coffee beans	1		
	Cranberry	0.05	(*)	
	Cucumber	1		

Edible offal (mammalian)	0.3		
Egg plant	0.2		
Eggs	0.02		
Grapes	1		
Hops, dry	10		
Kale (Collards, curly kale, Scotch kale, thousand-headed kale; not including Marrow-stem kele)	5		
Leek	0.05	(*)	
Lettuce, head	2		
Mango	0.2		
Meat (from mammals other than marine mammals)	0.1		
Melons, except watermelon	0.2		
Milks	0.1		
Olives for oil production	2		
Onion, bulb	0.1		
Peaches (including apricots and nectarine) (subgroup)	1.5		
Peanut	1		
Pear	1		
Peas (pods and succulent=immature seeds)	5		
Peas, shelled (succulent seeds)	2		
Peppers (subgroup)	1		
Peppers chili, dried	10		
Plums (including fresh prunes) (subgroup)	1.5		
Pomegranate	1		
Poultry meat	0.02		
Poultry, edible offal of	0.05		
Prunes	5		
Pulses (group)	2		(except soya beans)
Radish leaves (including radish tops)	5		
Rape seed	0.05	(*)	
Root and tuber vegetables (group)	0.5		
Soya bean (dry)	3		
Squash, summer	1		
Strawberry	0.5		
Sunflower seed	0.05	(*)	

	Sweet corn (corn-on-the-cob)	0.02	(*)	
	Table olives	2		
	Tea, green, black (black, fermented and dried)	50		
	Tomato	0.5		
	Tree nuts (group)	0.01		
	Watermelon	0.2		
	Wheat bran, unprocessed	0.3		
	Wheat flour	0.03		
	Wheat, hay and/or straw	1		
	Citrus (Acid Lime)	1		
	Groundnut Seed	1		
	Sugarcane	0.1		
	Okra	2		
	Sunflower Seed	0.5		
	Tomato	1		
	Cucumber	1		
	Cottonseed Oil	0.05		
	Rice	0.05		
	Brinjal	0.2		
	Soybean oil	0.01	*	
Indoxacarb	Apple	0.5		
	Broccoli	0.2		
	Cabbages, head	3		
	Cauliflower	0.2		
	Chick-pea (dry)	0.2		
	Cotton seed	1		
	Cowpea (dry)	0.1		
	Cranberry	1		
	Edible offal (mammalian)	0.05		
	Egg plant	0.5		
	Eggs	0.02		
	Fruiting vegetables, cucurbits (group)	0.5		
	Grape, dried (= Currants, Raisins and Sultanas)	5		
	Grapes	2		
	Lettuce, head	7		
	Lettuce, leaf	3		
	Meat (from mammals other than marine mammals)	2	(fat)	
	Milk fats	2		
	Milks	0.1		
	Mints	15		
	Mung bean (dry)	0.2		

	Peanut	0.02	(*)	
	Pear	0.2		
	Peppers (subgroup)	0.3		
	Potato	0.02		
	Poultry meat	0.01	(*) (fat)	
	Poultry, edible offal of	0.01	(*)	
	Prunes	3		
	Soya bean (dry)	0.5		
	Stone fruits (group)	1		
	Sweet corn (corn-on-the-cob)	0.02		
	Tea, green, black (black, fermented and dried)	5		
	Tomato	0.5		
	Pigeon pea	0.1		
	Rice	0.05		
	Cottonseed	1		
	Cottonseed Oil	0.1		
	Cabbage	3		
Iprodione	Almonds	0.2		
	Barley	2		
	Beans (dry)	0.1		
	Blackberries	30		
	Broccoli	25		
	Carrot	10	Po	
	Cherries (subgroup)	10		
	Common bean (pods and/or immature seeds)	2		
	Cucumber	2		
	Grapes	10		
	Kiwifruit	5		
	Lettuce, head	10		
	Lettuce, leaf	25		
	Onion, bulb	0.2		
	Peach	10		
	Pome fruits (group)	5	Po	
	Rape seed	0.5		
	Raspberries, red, black	30		
	Rice, husked	10		
	Spices, roots and rhizomes	0.1		
	Spices, seeds	0.05	(*)	
	Strawberry	10		
	Sugar beet	0.1	(*)	
	Sunflower seed	0.5		
	Tomato	5		
	Witloof chicory (sprouts)	1		

	Rape seed	0.5		
	Mustard seed	0.5		
Isoprothiolane	Edible offal (mammalian)	0.01	(*)	
	Mammalian fats (except milk fats)	0.01	(*)	
	Meat (from mammals other than marine mammals)	0.01	(*)	
	Milks	0.01	(*)	
	Rice, husked	6		
	Rice, polished	1.5		
Isoproturon	Wheat	0.1		
Kasugamycin	Rice	0.05		
	Tomato	0.05		
Kresoxim-Methyl	Barley, similar grains, and pseudocereals with husks (subgroup)	0.15		
	Beetroot	0.05	(*)	
	Currants, black, red, white	0.9		
	Edible offal (mammalian)	0.05		
	Eggs	0.02	(*)	
	Fruiting vegetables, cucurbits (group)	0.5		
	Garlic	0.01		
	Grape, dried (= Currants, Raisins and Sultanas)	3		
	Grapes	1.5		
	Leek	10		
	Mammalian fats (except milk fats)	0.02	(*)	
	Mango	0.1		
	Meat (from mammals other than marine mammals)	0.02	(*)	
	Milks	0.02	(*)	
	Olive oil, virgin	1		
	Olives for oil production	0.2		
	Peach	1.5		
	Pecan	0.05	(*)	
	Peppers, sweet (including pimento or pimienta)	0.3		
	Pome fruits (group)	0.15		(except Japanese persimmon)
	Poultry fats	0.02	(*)	
	Poultry meat	0.02	(*)	
	Poultry, edible offal of	0.02	(DM)	

	Straw and hay of cereal grains except pseudocereals	3	(*)	
	Sugar beet	0.05		
	Table olives	0.2		
	Turnip	0.05		
	Wheat, similar grains, and pseudocereals without husks (subgroup)	0.05		
	Meats	0.05		
	Maize	0.02	*	
	Chilli	0.15		
	Dried Chilli	1.5		
	Potato	0.02	*	
	Soybean	0.02	*	
	Soybean oil	0.02	*	
	soybean cake	0.02	*	
	Cotton seed Oil	0.02	*	
Lambda cyhalothrin	Brinjal	0.2		
	Tomato	0.1		
	Rice	1		
	Okra	2		
	Red Gram	0.05		
	Bengal Gram	0.05		
	Chilli	0.05		
	Dried Chilli	0.5		
	Groundnut seed	0.01		
	Onion	0.01		
	Soybean	0.01		
	Mango	0.2		
	Grapes	0.05		
	Tea	0.05	*	
	Maize	0.01	*	
	Cottonseed Oil	0.05		
Lufenuron	Coffee beans	0.07		
	Cucumber	0.09		
	Edible offal (mammalian)	0.15		
	Eggs	0.02		
	Lime	0.4		
	Maize	0.01		
	Mammalian fats (except milk fats)	2		
	Meat (from mammals other than marine mammals)	2	(fat)	
	Melons, except watermelon	0.4		
	Milk fats	5		



	Milks	0.15		
	Orange oil, edible	8		
	Oranges, sweet, sour (including Orange-like hybrids) (subgroup)	0.3		
	Peppers, sweet (including pimento or pimienta)	0.8		
	Pome fruits (group)	1		
	Potato	0.01	(*)	
	Poultry fats	0.04		
	Poultry meat	0.02		
	Poultry, edible offal of	0.02	(fat)	
	Soya bean (dry)	0.01	(*)	
	Tomato	0.4		
	Cauliflower	0.1		
	Cottonseed	0.01		
	Black gram	0.02	*	
	Chilli	0.05		
	Dried Chilli	0.5		
	Cabbage	0.3		
	Pigeon pea	0.01		
Malathion (Malathion to be determined and expressed as combined residues of malathion and malaoxon)	Apple	0.5		
	Asparagus	1		
	Beans (dry)	2		
	Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	1		
	Blueberries	10		
	Cherries (subgroup)	3		
	Citrus fruits (group)	7		
	Cotton seed	20		

	Cotton seed oil, crude	13		
	Cotton seed oil, edible	13		
	Cucumber	0.2		
	Grapes	5		
	Maize	0.05		
	Mustard greens	2		
	Onion, bulb	1		
	Peppers (subgroup)	0.1		
	Peppers chili, dried	1		
	Sorghum grain	3		
	Spices, fruits and berries	1		
	Spices, roots and rhizomes	0.5		
	Spices, seeds	2		
	Spinach	3		
	Spring onion	5		



	Strawberry	1		
	Sweet corn (corn-on-the-cob)	0.02		
	Tomato	0.5		
	Tomato juice	0.01		
	Turnip greens	5		
	Turnip, Garden	0.2		
	Wheat	10		
	Wheat bran, unprocessed	25		
	Wheat flour	0.2		
	Other food grains	4		
	Maize	0.05		
	Milled food grains	1		
	Fruits	4		
	Vegetables	3		
	Dried fruits	8		
	Carbonated Water	0.01		
Mancozeb	Chilli	1		
	Dried Chilli	10		
	Cauliflower	0.02		
	Groundnut	0.1		
	Cumin	10		
	Black Pepper	2		
	Mustard seed	0.1		
	Gherkin	0.1	*	
	Tea	3		
	Onion	4		
	Milks	0.05		
	Meats	0.1		
	Mango	2		
	Grapes	5		
	citrus	0.05	*	

	cucumber	0.4		
	rice	0.5	*	
Mandipropamid	Beans with pods (subgroup)	1		
	Broccoli	2		
	Cabbages, head	3		
	Cacao beans	0.06		
	Celery	20		
	Cucumber	0.2		
	Edible offal (mammalian)	0.01	(*)	
	Eggs	0.01	(*)	
	Grape, dried (= Currants, Raisins and Sultanas)	5		
	Grapes	2		
	Hops, dry	90		
	Leafy vegetables	25		
	Mammalian fats (except milk fats)	0.01	(*)	
	Meat (from mammals other than marine mammals)	0.01	(*)	
	Melons, except watermelon	0.5		
	Milks	0.01	(*)	
	Onion, bulb	0.1		
	Peppers (subgroup)	1		
	Peppers chili, dried	10		
	Potato	0.1		
	Poultry fats	0.01	(*)	
	Poultry meat	0.01	(*)	
	Poultry, edible offal of	0.01	(*)	
	Spring onion	7		
	Squash, summer	0.2		
	Tomato	0.3		
Methyl Chlorophenoxy Acetic Acid (MCPA)	Barley	0.2		
	Barley, hay and/or straw	50		
	Edible offal (mammalian)	3		
	Eggs	0.05	(*)	
	Flax-seed	0.01	(*)	
	Maize	0.01	(*)	
	Mammalian fats (except milk fats)	0.2		
	Meat (from mammals other than marine mammals)	0.1		
	Milks	0.04		
	Oats	0.2		
	Peas (dry)	0.01	(*)	

	Poultry fats	0.05	(*)	
	Poultry meat	0.05	(*)	
	Poultry, edible offal of	0.05	(*)	
	Rye	0.2		
	Triticale	0.2		
	Triticale, hay and/or straw	50		
	Wheat	0.2		
	Wheat, hay and/or straw	50		
	Rice	0.05		
Mesotrione	Almond hulls	0.04	(dw)	
	Asparagus	0.01	(*)	
	Bush berries (subgroup)	0.01	(*)	
	Cane berries (subgroup)	0.01	(*)	
	Citrus fruits (group)	0.01	(*)	
	Cranberry	0.01	(*)	
	Edible offal (mammalian)	0.01	(*)	
	Eggs	0.01	(*)	
	Linseed	0.01	(*)	
	Maize	0.01	(*)	
	Meat (from mammals other than marine mammals)	0.01	(*)	
	Milks	0.01	(*)	
	Millet (Including barnyard, bulrush, common, finger, foxtail, little Millet)	0.01	(*)	
	Oats	0.01	(*)	
	Okra	0.01	(*)	
	Pome fruits (group)	0.01	(*)	
	Poultry meat	0.01	(*)	
	Poultry, edible offal of	0.01	(*)	
	Rhubarb	0.01	(*)	
	Rice, husked	0.01	(*)	
	Sorghum grain	0.01	(*)	
	Soya bean (dry)	0.03		
	Stone fruits (group)	0.01	(*)	
	Sugar cane	0.01		
	Sweet corn (corn-on-the-cob)	0.01	(*)	
	Tree nuts (group)	0.01	(*)	
Metalaxyl	Asparagus	0.05	(*)	
	Avocado	0.2		
	Broccoli	0.5		
	Brussels sprouts	0.2		
	Cabbages, head	0.5		
	Cacao beans	0.2		
	Carrot	0.05	(*)	

	Cauliflower	0.5		
	Cereal grains	0.05	(*)	
	Citrus fruits (group)	5	Po	
	Cotton seed	0.05	(*)	
	Cucumber	0.5		
	Gherkin	0.5		
	Grapes	1		
	Hops, dry	10		
	Lettuce, head	2		
	Melons, except watermelon	0.2		Withdrawal recommended by JMPR (2021)
	Onion, bulb	2		
	Peanut	0.1		
	Peas, shelled (succulent seeds)	0.05	(*)	
	Peppers (subgroup)	1		
	Peppers chili, dried	10		
	Pome fruits (group)	1	Po	
	Potato	0.05	(*)	
	Raspberries, red, black	0.2		
	Soya bean (dry)	0.05	(*)	
	Spices, seeds	5		
	Spinach	2		
	Squash, summer	0.2		
	Sugar beet	0.05	(*)	
	Sunflower seed	0.05	(*)	
	Tomato	0.5		
	Watermelon	0.2		
	Winter squash	0.2		
	Pearl Millet (Bajra)	0.05		
	Maize	0.05		
	Sorghum	0.05		
Metalaxyl-M	Potato	0.05	*	
	Grapes	1		
	Black pepper	0.5		
	Mustard Seed	0.01		
	Dried Chilli	0.2		
	Tomato	0.5		
	Chilli	0.02		
Methoxyfenozide	Almond hulls	50		
	Avocado	0.7		
	Beans (dry)	0.5		
	Beans, shelled	0.3		
	Blueberries	4		
	Broccoli	3		

Cabbages, head	7		
Carrot	0.5		
Celery	15		
Citrus fruits (group)	2		
Common bean (pods and/or immature seeds)	2		
Cotton seed	7		
Cowpea (dry)	5		
Cranberry	0.7		
Edible offal (mammalian)	0.2		
Eggs	0.01		
Fruiting vegetables, cucurbits (group)	0.3		(except watermelon)
Grape, dried (= Currants, Raisins and Sultanas)	2		
Grapes	1		
Lettuce, head	15		
Lettuce, leaf	30		
Maize	0.02	(*)	
Mammalian fats (except milk fats)	0.3		
Meat (from mammals other than marine mammals)	0.3	(fat)	
Milks	0.05		
Mustard greens	30		
Papaya	1		
Peanut	0.03		
Peanut oil, edible	0.1		
Peas (dry)	5		
Peas (pods and succulent=immature seeds)	2		
Peas, shelled (succulent seeds)	0.3		
Peppers (subgroup)	2		
Peppers chili, dried	20		
Pome fruits (group)	2		
Poultry meat	0.01	(*)	
Poultry, edible offal of	0.01	(*)	
Prunes	2		
Radish	0.4		
Radish leaves (including radish tops)	7		
Stone fruits (group)	2		
Strawberry	2		
Sugar beet	0.3		

	Sweet corn (corn-on-the-cob)	0.02	(*)	
	Sweet potato	0.02		
	Tomato	2		
Methoxyfenozide	Tree nuts (group)	0.1		
Methyl Bromide	Bread and other cooked cereal products	0.01	(*)	To apply to commodity at point of retail sale or when offered for consumption
	Cacao beans	5	Po	To apply at point of entry into a country and, in case of cereal for milling, if product has been freely exposed to air for a period of at least 24 h after fumigation and before
	Cereal grains	5	Po	To apply at point of entry into a country and, in case of cereal for milling, if product has been freely exposed to air for a period of at least 24 h after fumigation and before
	Cocoa products	0.01	(*)Po	To apply to commodity at point of retail sale or when offered for consumption
	Dried fruits	2	Po	To apply at point of entry into a country and, in case of cereal for milling, if product has been freely exposed to air for a period of at least 24 h after fumigation and before
	Dried fruits	0.01	(*)Po	To apply to commodity at point of retail sale or when offered for consumption





	Milled cereals products	0.01	(*)Po	To apply to commodity at point of retail sale or when offered for consumption
	Milled cereals products	1	Po	To apply at point of entry into a country and, in case of cereal for milling, if product has been freely exposed to air for a period of at least 24 h after fumigation and before
	Peanut	0.01	(*)Po	To apply to commodity at point of retail sale or when offered for consumption
	Peanut	10	Po	To apply at point of entry into a country and, in case of cereal for milling, if product has been freely exposed to air for a period of at least 24 h after fumigation and before
	Tree nuts (group)	0.01	(*)Po	To apply to commodity at point of retail sale or when offered for consumption
	Tree nuts (group)	10	Po	To apply at point of entry into a country and, in case of cereal for milling, if product has been freely exposed to air for a period of at least 24 h after fumigation and before
Metiram as CS2	Chilli	1		
	Dry chilli	10		
	Grapes	5		
	Potato	0.2		
	Tomato	5		



	Groundnut seed	0.1		
	Groundnut seed oil	0.1		
	Milks	0.05		
	Onion	0.05	*	
	Apple	0.05	*	
	Cotton seed	0.05	*	
	Cotton seed oil	0.05	*	
	cumin	10		
	Banana	2		
	Black gram	0.05	*	
	cucumber	2		
	pomegranate	0.05	*	
	green gram	0.05	*	
Metolachlor	Soybean Oil	0.05		
Metribuzin	Milks	0.01	*	
	Tomato	0.05	*	
	Sugarcane	0.01	*	
	Potato	0.05	*	
	Soybean Oil	0.1		
Metsulfuron Methyl	Wheat	0.03		
	Rice	0.01		
	Wheat	0.1		
Novaluron	Sugarcane	0.02		
	Beans (dry)	0.1		
	Blueberries	7		
	Brassica vegetables (except Brassica leafy vegetables) (group)	0.7		
	Chard	15		
	Common bean (pods and/or immature seeds)	0.7		
	Cotton seed	0.5		
	Edible offal (mammalian)	0.7		
	Eggs	0.1		
	Fruiting vegetables, cucurbits (group)	0.2		
	Fruiting vegetables, other than cucurbits (group)	0.7		Except sweet corn.
	Meat (from mammals other than marine mammals)	10	(fat)	
	Milk fats	7		
	Milks	0.4		
	Mustard greens	25		
	Pome fruits (group)	3		
	Potato	0.01	(*)	
	Poultry meat	0.5	(fat)	

	Poultry, edible offal of	0.1		
	Prunes	3		
	Soya bean (immature seeds)	0.01	(*)	
	Stone fruits (group)	7		
	Strawberry	0.5		
	Sugar cane	0.5		
	Chilli	0.01		
	Dried Chilli	0.1		
	Cottonseed Oil	0.01		
	Cabbage	0.7		
Orthosulfamuron	Paddy	0.1		
Oxadiargyl	Mustard Seed	0.05		
	Onion	0.1		
	Cumin	0.01		
	Rice	0.1		
	Sunflower Seed	0.05	*	
	Sunflower oil	0.05	*	
Oxadiazon	Rice	0.03		
Oxyfluorfen	Rice	0.05		
	Groundnut Oil	0.05		
	Mentha	0.01		
	Tea	0.2		
	Potato	0.01		
	Onion	0.05		
Paraquat dichloride (Determined as Paraquatcations)	Almond hulls	0.01	(*)	
	Assorted tropical and sub-tropical fruits - inedible peel	0.01	(*)	
	Berries and other small fruits	0.01	(*)	
	Citrus fruits (group)	0.02		
	Cotton seed	2		
	Edible offal (mammalian)	0.05		
	Eggs	0.005	(*)	
	Fruiting vegetables, cucurbits (group)	0.02		
	Fruiting vegetables, other than cucurbits (group)	0.05		
	Hops, dry	0.1		

	Leafy vegetables	0.07		
	Maize	0.03		
	Maize flour	0.05		
	Meat (from mammals other than marine mammals)	0.005		
	Pome fruits (group)	0.01	(*)	
	Poultry meat	0.005	(*)	
	Poultry, edible offal of	0.005	(*)	
	Pulses (group)	0.5		
	Rice	0.05		
	Rice, hay and/or straw	0.05		
	Root and tuber vegetables (group)	0.05		
	Sorghum grain	0.03		
	Stone fruits (group)	0.01	(*)	
	Sunflower seed	2		
	Table olives	0.1		
	Tea, green, black (black, fermented and dried)	0.2		
	Tree nuts (group)	0.05		
	Other Food grains	0.1		
	Sorghum	0.03		
	Milled food grains	0.03		
	Potato	0.2		
	Other vegetables	0.05		
	Cottonseed	2		



	Cottonseed oil (edible refined)	0.05		
	Milks	0.01		
	Fruits	0.05		
Penconazole	Apple	0.1		
	Artichoke, globe	0.06		
	Cucumber	0.06		
	Currant, black	2		
	Edible offal (mammalian)	0.05	(*)	
	Egg plant	0.09		
	Eggs	0.05	(*)	
	Gherkin	0.06		
	Grape, dried (= Currants, Raisins and Sultanias)	1.5		
	Grapes	0.4		
	Mammalian fats (except milk fats)	0.05	(*)	
	Meat (from mammals other than marine mammals)	0.05	(*)	
	Melons, except watermelon	0.15		
	Milks	0.01	(*)	
	Peaches (including apricots and nectarine) (subgroup)	0.08		
	Pear	0.1		
	Peppers, sweet (including pimento or pimienta)	0.2		
	Poultry meat	0.05	(*)	
	Poultry, edible offal of	0.05	(*)	
	Squash, summer	0.06		
	Strawberry	0.5		
	Tomato	0.09		
	Black gram	0.02		
	Mango	0.05		
	Apple	0.1		
Pendimethalin	Almond hulls	7	(dw)	
	Asparagus	0.1		
	Beans (dry)	0.05		
	Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	0.05		(green pods and immature seeds)
	Bush berries (subgroup)	0.05	(*)	
	Cane berries (subgroup)	0.05	(*)	

	Carrot	0.5		
	Celery	0.09		
	Citrus fruits (group)	0.03		
	Edible offal (mammalian)	0.05		
	Eggs	0.01	(*)	
	Fennel, bulb	0.05	(*)	
	Garlic	0.05	(*)	
	Hops, dry	0.05		
	Kale (Collards, curly kale, Scotch kale, thousand-headed kale; not including Marrow-stem kele)	0.5		
	Leaves of Brassicaceae	0.3		except kale
	Lettuce, leaf	4		
	Mammalian fats (except milk fats)	0.2		
	Meat (from mammals other than marine mammals)	0.2	(fat)	
	Milk fats	0.8		
	Milks	0.02		
	Mints	0.2		
	Onion, bulb	0.05	(*)	
	Onion, Welsh	0.4		
	Peas (dry)	0.05		
	Peas (pods and succulent=immature seeds)	0.05		
	Peas, shelled (succulent seeds)	0.05		
	Peppermint oil, edible	6		
	Poultry fats	0.01	(*)	
	Poultry meat	0.01	(*)	
	Poultry, edible offal of	0.01	(*)	
	Shallot	0.05	(*)	
	Spring onion	0.4		
	Strawberry	0.05	(*)	
	Tree nuts (group)	0.05		
	Wheat	0.05		
	Rice	0.05		
	Soybean Oil	0.05		
	Cottonseed Oil	0.05		
	Chilli	0.05	*	
	Dried Chilli	0.5		
	Red Gram	0.05	*	
Penoxuslum	Rice	0.1	*	
Phenthoate	Food grains	0.05		
	Milled food grains	0.01		







	Oil seeds	0.03		
	Edible oils	0.01		
	Eggs	0.05		(shell free basis)
	Meat & Poultry	0.05		(Carcass fat basis)
	Milks	0.01		(F)
	Spices, seeds	7		
Phosmet	Apricot	10		
	Blueberries	10		
	Cattle meat	1	(fat) V	
	Citrus fruits (group)	3		
	Cotton seed	0.05		
	Cranberry	3		
	Grapes	10		
	Milks	0.02		
	Nectarine	10		
	Peach	10		
	Pome fruits (group)	10		
	Potato	0.05	(*)	
	Tree nuts (group)	0.2		
Pirimiphos-Methyl	Cereal grains	7	Po	
	Edible offal (mammalian)	0.01	(*)	
	Spices, fruits and berries	0.5		
	Spices, seeds	3		
	Wheat bran, unprocessed	15		
	Rice	0.5		
	Milled food grains except rice	1		
	Eggs	0.05		(shell free basis)
	Meats & Poultry	0.05		(Carcass fat basis)
	Milks	0.05		(F)
Pretilachlor	Rice	0.05		
Profenofos	Cardamom	3		
	Coffee beans	0.04		
	Coriander, seed	0.1		
	Cotton seed	3		
	Cumin seed	5		
	Edible offal (mammalian)	0.05	(*)	
	Eggs	0.02	(*)	
	Fennel, seed	0.1		
	Mango	0.2		
	Mangostan	10		

	Meat (from mammals other than marine mammals)	0.05	(*)	
	Milks	0.01	(*)	
	Peppers chili	3		
	Peppers chili, dried	20		
	Poultry meat	0.05	(*)	
	Poultry, edible offal of	0.05	(*)	
	Spices, fruits and berries	0.07		(except cardamom)
	Spices, roots and rhizomes	0.05		
	Teas (tea and herb teas)	0.5		
	Tomato	10		
	Cottonseed oil	3		
	Soybean	0.01	*	
Propamocarb	Broccoli	3		
	Brussels sprouts	2		
	Cabbages, head	1		
	Cauliflower	2		
	Edible offal (mammalian)	1.5		
	Egg plant	0.3		
	Eggs	0.01	(*)	
	Fruiting vegetables, cucurbits (group)	5		
	Kale (Collards, curly kale, Scotch kale, thousand-headed kale; not including Marrow-stem kale)	20		
	Leek	30		
	Lettuce, head	100		
	Lettuce, leaf	100		
	Mammalian fats (except milk fats)	0.03		
	Meat (from mammals other than marine mammals)	0.03		
	Milks	0.01	(*)	
	Onion, bulb	2		
	Peppers chili, dried	10		
	Peppers, sweet (including pimento or pimienta)	3		
	Potato	0.3		
	Poultry fats	0.01	(*)	
	Poultry meat	0.01	(*)	
	Poultry, edible offal of	0.01	(*)	
	Radish	1		
	Spinach	40		
	Tomato	2		

	Witloof chicory (sprouts)	2		
Propanil	Rice	0.05	*	
Propargite	Almond hulls	50		
	Almonds	0.1	(*)	
	Apple	3		
	Apple juice	0.2		
	Beans (dry)	0.3		
	Broad bean (dry)	0.3		
	Chick-pea (dry)	0.3		
	Citrus fruits (group)	3		
	Citrus pulp, dried	10		
	Cotton seed	0.1		
	Cotton seed oil, edible	0.2		
	Edible offal (mammalian)	0.1	(*)	
	Eggs	0.1	(*)	
	Grape juice	1		
	Grape pomace, dry	40		
	Grape, dried (= Currants, Raisins and Sultanas)	12		
	Grapes	7		
	Hops, dry	100		
	Lipin (dry)	0.3		
	Maize	0.1	(*)	
	Maize flour	0.2		
	Maize oil, crude	0.7		
	Maize oil, edible	0.5		
	Meat (from mammals other than marine mammals)	0.1	(*) (fat)	
	Milks	0.1	(*) F	
	Orange juice	0.3		
	Peanut	0.1	(*)	
	Peanut oil, crude	0.3		
	Peanut oil, edible	0.3		
	Potato	0.03		
	Poultry meat	0.1	(*) (fat)	
	Poultry, edible offal of	0.1	(*)	
	Stone fruits (group)	4		
	Tomato	2		
	Walnuts	0.3		
	Brinjal	2		
	Chilli	2		
	Dried Chilli	20		
	Apple	3		
	Tea	10		
Propiconazole	Banana	0.1		

Barley	2		
Barley, hay and/or straw	8		
Cherries (subgroup)	3	Po	
Coffee beans	0.02		
Cranberry	0.3		
Edible offal (mammalian)	0.5		
Eggs	0.01	(*)	
Lemons and limes (including citron) (subgroup)	10	Po	
Maize	0.05		
Mammalian fats (except milk fats)	0.01	(*)	
Mandarins (including mandarin-like hybrids) (subgroup)	10	Po	
Meat (from mammals other than marine mammals)	0.01	(*) (fat)	
Milks	0.01	(*)	
Oats	0.7		
Orange oil, edible	1850		
Oranges, sweet, sour (including Orange-like hybrids) (subgroup)	10	Po	
Peaches (including apricots and nectarine) (subgroup)	4	Po	
Pecan	0.02	(*)	
Pineapple	2	Po	
Plums (including fresh prunes) (subgroup)	0.4	Po	
Popcorn	0.05		
Poultry meat	0.01	(*) (fat)	
Pummelo and grapefruits (including Shaddock-like hybrids, among others Grapefruit) (subgroup)	4	Po	
Rape seed	0.02		
Rye	0.09		
Soya bean (dry)	0.07		
Sugar beet	0.02		
Sugar cane	0.02	(*)	
Sweet corn (corn-on-the- cob)	0.05		
Tomato	3	Po	
Triticale	0.09		
Triticale, hay and/or straw	15		
Wheat	0.09		



	Wheat, hay and/or straw	15		
	Tea	0.1		
	Groundnut seed	0.1		
	Rice	0.05		
Propineb	Rice	0.05		
	Tomato	1		
	Apple	1		
	Pomegranate	0.5		
	Potato	0.5		
	Chilli	2		
	Dried Chilli	20		
	Grapes	0.5		
Pymetrozine	Acerola	undefined		
	Rice	0.01	*	
Pyraclostrobin	Artichoke, globe	2		
	Asparagus	0.01	(*)	
	Avocado	0.2		
	Banana	0.02	(*)	
	Barley	1		
	Beans (dry)	0.2		
	Beans with pods (subgroup)	0.3		(except common beans)
	Blackberries	3		
	Blueberries	4		
	Broad bean, without pods (succulent seeds)	0.01		
	Brussels sprouts	0.3		
	Cabbages, head	0.2		
	Cacao beans	0.01		
	Cantaloupe	0.2		
	Celery	1.5		
	Cherries (subgroup)	3		(includes all commodities in this subgroup)
	Citrus fruits (group)	2		
	Citrus oil, edible	10		
	Coffee beans	0.3		
	Common bean (pods and/or immature seeds)	0.6		
	Common beans (succulent seeds)	0.3		
	Dry peas (subgroup)	0.3		
	Edible offal (mammalian)	0.05		
	Egg plant	0.3		
	Eggs	0.05	(*)	

Flowerhead brassicas (includes Broccoli: Broccoli, Chinese and cauliflower)	0.1		
Fruiting vegetables, cucurbits (group)	0.5		
Garlic	0.15		
Grape, dried (= Currants, Raisins and Sultanas)	5		
Grapes	2		
Hops, dry	15		
Kale (Collards, curly kale, Scotch kale, thousand- headed kale; not including Marrow-stem kele)	1		
Leek	0.7		
Lentil (dry)	0.5		
Lettuce, head	40		
Maize	0.02	(*)	
Mammalian fats (except milk fats)	0.5		
Mango	0.6		
Meat (from mammals other than marine mammals)	0.5	(fat)	
Milks	0.03		
Oats	1		
Oilseed, except peanut	0.4		
Olive oil, virgin	0.07		
Olives for oil production	0.01		
Onion, bulb	1.5		
Papaya	0.15		
Passion fruit	0.2		
Peaches (including apricots and nectarine) (subgroup)	0.3		
Peanut, whole	0.02	(*)	
Peas with pods (subgroup)	0.3		
Peppers (subgroup)	0.5		
Pineapple	0.3		
Pistachio nuts	1		
Plums (including fresh prunes) (subgroup)	0.8		(includes all commodities in this subgroup)
Pome fruits (group)	0.7		
Poultry meat	0.05	(*) (fat)	
Poultry, edible offal of	0.05	(*)	

	Radish leaves (including radish tops)	20		
	Raspberries, red, black	3		
	Rice	1.5		
	Rice, hay and/or straw	5	(dw)	
	Rice, husked	0.09		
	Rice, polished	0.03		
	Root vegetables (subgroup)	0.5		(except sugar beet)
	Rye	0.2		
	Sorghum grain	0.5		
	Soya bean (dry)	0.05		
	Spelt	0.2		
	Spinach	0.6		
	Spring onion	1.5		
	Straw and hay of cereal grains except pseudocereals	30		
	Strawberry	1.5		
	Succulent peas without pods (subgroup)	0.08		
	Sugar cane	0.08		
	Table olives	0.01		
	Tea, green, black (black, fermented and dried)	6		
	Tomato	0.3		
	Tree nuts (group)	0.02	(*)	except pistachio nuts
	Triticale	0.2		
	Tuberous and corm vegetables (subgroup)	0.02	(*)	
	Wheat	0.2		
	Witloof chicory (sprouts)	0.09		
	Cumin	0.02	*	
	Banana	0.02	*	
	Pomegranate	0.02	*	
	Potato	0.05	*	
Pyrazosulfuron ethyl	Rice	0.01		
Pyridalyl	Cottonseed Oil	0.02		
	Cabbage	0.02		
	Okra	0.02		
	Dried Chilli	0.2		
	Chilli	0.02		
Pyriproxyfen	Cattle meat	0.01	(*) (fat)	
	Cattle, edible offal of	0.01	(*)	
	Citrus fruits (group)	0.5		
	Cotton seed	0.05		

	Cotton seed oil, crude	0.01		
	Cotton seed oil, edible	0.01		
	Cucumber	0.04		
	Egg plant	0.6		
	Gherkin	0.04		
	Goat meat	0.01	(*) (fat)	
	Goat, edible offal of	0.01	(*)	
	Mango	0.02	(*)	
	Melons, except watermelon	0.07		
	Papaya	0.3		
	Peppers (subgroup)	0.6		
	Peppers chili, dried	6		
	Pineapple	0.01		
	Squash, summer	0.04		
	Tomato	0.4		
	Brinjal	0.02		
	Okra	0.03	*	
Quinalphos	Cauliflower	0.1		
	Citrus	0.05		
	Bengal Gram	0.05		
	Cottonseed Oil	0.05		
	Mustard seed oil	0.1		
	Soybean	0.05		
	Groundnut oil	0.3		
	Rice	0.01		
	Pigeon pea	0.01		
	Cardamom	0.01		
	Tea	0.01		
	Fish	0.01		
	Chilli	0.2		
	Dried Chilli	2		
Quinclorac	Cranberry	1.5		
	Edible offal (mammalian)	0.1		
	Eggs	0.05	(*)	
	Mammalian fats (except milk fats)	0.05	(*)	
	Meat (from mammals other than marine mammals)	0.05	(*) (fat)	
	Milks	0.05	(*)	
	Poultry fats	0.05	(*)	
	Poultry meat	0.05		
	Poultry, edible offal of	0.05	(*)	
	Rape seed	0.15		
	Rhubarb	0.5		
	Rice	10		



	Rice, hay and/or straw	8	(dw)	
	Rice, husked	10		
	Rice, polished	8		
Quizalofop ethyl	Cottonseed	0.1		
	Soybean seed	0.05		
	Onion	0.01	*	
	Groundnut	0.1		
	Black gram	0.01	*	
Quizalofop-P-tefuryl	Soybean Seed	0.02		
	Cotton seed	0.05	*	
	Cotton seed oil	0.05	*	
Spinosad	Apple	0.1		
	Blackberries	1		
	Blueberries	0.4		
	Brassica vegetables (except Brassica leafy vegetables) (group)	2		
	Cattle kidney	1		The MRL accommodates external animal treatment.
	Cattle liver	2		The MRL accommodates external animal treatment.
	Cattle meat	3	(fat)	The MRL accommodates external animal treatment.
	Cattle milk	1		The MRL accommodates external animal treatment.
	Cattle milk fat	5		
	Celery	2		
	Cereal grains	1	Po	
	Citrus fruits (group)	0.3		
	Cotton seed	0.01	(*)	
	Cotton seed oil, crude	0.01	(*)	
	Cotton seed oil, edible	0.01	(*)	
	Cranberry	0.02		
	Dewberries (including boysenberry and loganberry)	1		
	Edible offal (mammalian)	0.5		Except cattle
	Eggs	0.01		

	Fruiting vegetables, cucurbits (group)	0.2		
	Grape, dried (= Currants, Raisins and Sultanas)	1		
	Grapes	0.5		
	Kiwifruit	0.05		
	Leafy vegetables	10		
	Legume vegetables (group)	0.3		
	Meat (from mammals other than marine mammals)	2	(fat)	
	Onion, bulb	0.1		
	Passion fruit	0.7		
	Peppers (subgroup)	0.3		
	Peppers chili, dried	3		
	Potato	0.01	(*)	
	Poultry meat	0.2	(fat)	
	Raspberries, red, black	1		
	Soya bean (dry)	0.01	(*)	
	Spring onion	4		
	Stone fruits (group)	0.2		
	Sweet corn (corn-on-the-cob)	0.01	(*)	
	Tomato	0.3		
	Tree nuts (group)	0.07		
	Wheat bran, unprocessed	2		
	Wheat, hay and/or straw	1		
	Cabbage	2		
	Red gram	0.01		
Spirodiclofen	Almond hulls	15		
	Avocado	0.9		
	Blueberries	4		
	Citrus fruits (group)	0.4		
	Coffee beans	0.03	(*)	
	Cucumber	0.07		
	Currants, black, red, white	1		
	Edible offal (mammalian)	0.05	(*)	
	Gherkin	0.07		
	Grape, dried (= Currants, Raisins and Sultanas)	0.3		
	Grapes	0.2		
	Hops, dry	40		
	Meat (from mammals other than marine mammals)	0.01	(*) (fat)	
	Milks	0.004	(*)	
	Papaya	0.03	(*)	

Peppers, sweet (including pimento or pimienta)	0.2		
Pome fruits (group)	0.8		
Stone fruits (group)	2		
Strawberry	2		
Tomato	0.5		
Tree nuts (group)	0.05		
Brassica vegetables (except Brassica leafy vegetables) (group)	3		
Cassava	0.02	(*)	
Coffee beans	0.2		
Common bean (pods and/or immature seeds)	1		
Cotton seed	0.7		
Cucumber	0.15		
Edible offal (mammalian)	0.3		
Egg plant	0.7		
Eggs	0.02		
Fruiting vegetables, cucurbits (group)	0.09		except melon and cucumber
Leafy vegetables	15		
Leaves of Brassicaceae	15		
Low growing berries (subgroup)	3		
Maize	0.02	(*)	
Mammalian fats (except milk fats)	0.15		
Meat (from mammals other than marine mammals)	0.15	F	
Melons, except watermelon	0.3		
Milks	0.01		
Okra	0.5		
Pepino	0.5		
Peppers (subgroup)	0.5		
Peppers chili, dried	5		
Popcorn	0.02	(*)	
Potato	0.02	(*)	
Poultry fats	0.02		
Poultry meat	0.02		
Poultry, edible offal of	0.05		
Sweet corn (corn-on-the-cob)	0.02	(*)	
Sweet potato	0.02	(*)	
Tea, green, black (black, fermented and dried)	70		



	Tomato	0.7		
	Tomato paste	2		
	Tomato, dried	4		
Spirotetramat	Almond hulls	10		
	Artichoke, globe	1		
	Avocado	0.4		
	Bush berries (subgroup)	1.5		
	Cabbages, head	2		
	Carrot	0.04		
	Celery	4		
	Citrus fruits (group)	0.5		
	Cotton seed	0.4		
	Cotton seed, meal	1		
	Cranberry	0.2		
	Edible offal (mammalian)	1		
	Eggs	0.01		
	Flowerhead brassicas (includes Broccoli: Broccoli, Chinese and cauliflower)	1		
	Fruiting vegetables, cucurbits (group)	0.2		
	Fruiting vegetables, other than cucurbits (group)	1		Except sweet corn, mushrooms and chili pepper
	Grape pomace, dry	4		
	Grape, dried (= Currants, Raisins and Sultanas)	4		
	Grapes	2		
	Guava	2		
	Hops, dry	15		
	Kiwifruit	0.02	(*)	
	Leafy vegetables	7		
	Legume vegetables (group)	1.5		
	Litchi	15		
	Mango	0.3		
	Meat (from mammals other than marine mammals)	0.05		
	Milks	0.005		
	Onion, bulb	0.4		
	Papaya	0.4		
	Peppers chili	2		Non-bell
	Peppers chili, dried	20		
	Pome fruits (group)	0.7		
	Potato	0.8		

	Poultry meat	0.01	(*)	
	Poultry, edible offal of	0.01		
	Prunes	5		
	Pulses (group)	2		except soya bean (dry)
	Soya bean (dry)	4		
	Stone fruits (group)	3		
	Strawberry	0.3		
	Sugar beet	0.06		
	Sugar beet leaves or tops (dry)	8	(dw)	
	Sugar beet molasses	0.3		
	Sweet corn	1.5		
	Tree nuts (group)	0.5		
Sulfentrazone and its metabolite Desmethylsulfentrazone and 3-Hydroxymethylsulfentrazone	Soybean	0.2		
	Soybean oil	0.2		
	Soybean deoiled cake	0.2		
Sulfoxaflo	Barley	0.6		
	Barley, hay and/or straw	3		
	Beans (dry)	0.3		
	Broccoli	3		
	Cabbages, head	0.4		
	Carrot	0.05		
	Cauliflower	0.04		
	Celery	1.5		
	Cherries (subgroup)	1.5		
	Cotton seed	0.4		
	Cotton seed oil	0.4		
	Edible offal (mammalian)	1		
	Eggs	0.1		
	Fruiting vegetables, cucurbits (group)	0.5		
	Fruiting vegetables, other than cucurbits (group)	1.5		(except sweet corn and mushrooms)
	Garlic	0.01	(*)	
	Grape, dried (= Currants, Raisins and Sultanas)	6		
	Grapes	2		
	Leafy vegetables	6		

Lemons and limes (including citron) (subgroup)	0.4		
Maize	0.01	(*)	
Mammalian fats (except milk fats)	0.2		
Mandarins (including mandarin-like hybrids) (subgroup)	0.8		
Meat (from mammals other than marine mammals)	0.4		
Milks	0.3		
Onion, bulb	0.01	(*)	
Oranges, sweet, sour (including Orange-like hybrids) (subgroup)	0.8		
Peaches (including apricots and nectarine) (subgroup)	0.4		
Peppers chili, dried	15		
Plums (including fresh prunes) (subgroup)	0.5		
Pome fruits (group)	0.3		
Poultry fats	0.03		
Poultry meat	0.7		
Poultry, edible offal of	0.3		
Pummelo and grapefruits (including Shaddock-like hybrids, among others Grapefruit) (subgroup)	0.15		
Rape seed	0.15		
Rice	7		
Rice, hay and/or straw	20		
Rice, husked	1.5		
Rice, polished	1		
Root and tuber vegetables (group)	0.03		(except carrots)
Sorghum grain	0.2		
Soya bean (dry)	0.3		
Spring onion	0.7		
Strawberry	0.5		
Sweet corn (corn on the cob) (kernels plus cob with husk removed)	0.01	(*)	
Tree nuts (group)	0.03		
Triticale	0.2		
Wheat	0.2		
Wheat, hay and/or straw	3		



Tebuconazole	Apple	1		
	Apricot	2		
	Artichoke, globe	0.6		
	Asparagus	0.02	(*)	
	Banana	1.5		
	Barley	2		
	Barley, hay and/or straw	40		
	Beans (dry)	0.3		
	Beans with pods (subgroup)	3		
	Black gram	0.01	*	
	Broccoli	0.2		
	Brussels sprouts	0.3		
	Cabbage	1		
	Cabbages, head	1		
	Carrot	0.4		
	Cauliflower	0.05	(*)	
	Cherries (subgroup)	4		
	Citrus pulp, dried	3	(dw)	
	Coffee beans	0.1		
	Cotton seed	2		
	Cotton seed Oil	2		
	Cucumber	0.2		
	Edible offal (mammalian)	0.2		
	Egg plant	0.1		
	Eggs	0.05	(*)	
	Elderberries	1.5		
	Garlic	0.1		
	Ginseng	0.15		
	Ginseng, dried including red ginseng	0.4		
	Ginseng, extracts	0.5		
	Grape, dried (= Currants, Raisins and Sultanas)	7		
	Grapes	6		
	Hops, dry	40		
	Leek	0.7		
	Lettuce, head	5		
	Mandarins (including mandarin-like hybrids) (subgroup)	0.7	Po	
	Mango	0.2		
	Maize	0.05	*	
	Meat (from mammals other than marine mammals)	0.05	(*)	
	Melons, except watermelon	0.15		

	Milks	0.01	(*)	
	Nectarine	2		
	Oats	2		
	Onion, bulb	0.15		
	Orange oil, edible	10		
	Oranges, sweet, sour (including Orange-like hybrids) (subgroup)	0.4	Po	
	Papaya	2		
	Passion fruit	0.1		
	Peach	2		
	Peanut	0.15		
	Pear	1		
	Peppers chili, dried	10		
	Peppers, sweet (including pimento or pimiento)	1		
	Plums (including fresh prunes) (subgroup)	1		except prunes
	Poultry meat	0.05	(*)	
	Poultry, edible offal of	0.05	(*)	
	Prunes	3		
	Rape seed	0.3		
	Rice	1.5		
	Rye	0.15		
	Shallot	0.15		
	Soya bean (dry)	0.15		
	Spring onion	2		
	Squash, summer	0.2		
	Sunflower seed	0.1		
	Sweet corn (corn-on-the- cob)	0.6		
	Table olives	0.05	(*)	
	Tomato	0.7		
	Tree nuts (group)	0.05	(*)	
	Triticale	0.15		
	Wheat	0.15		
	Wheat, hay and/or straw	40		
Tebufenozide	Almond hulls	30		
	Almonds	0.05		
	Avocado	1		
	Blueberries	3		
	Broccoli	0.5		
	Cabbages, head	5		
	Cattle milk	0.05		
	Citrus fruits (group)	2		
	Cranberry	0.5		
	Edible offal (mammalian)	0.02	(*)	



	Eggs	0.02	(*)	
	Grape, dried (= Currants, Raisins and Sultanas)	2		
	Grapes	2		
	Kiwifruit	0.5		
	Leafy vegetables	10		
	Meat (from mammals other than marine mammals)	0.05	(fat)	
	Milks	0.01	(*)	
	Mints	20		
	Nectarine	0.5		
	Peach	0.5		
	Pecan	0.01	(*)	
	Peppers (subgroup)	1		
	Peppers chili, dried	10		
	Pome fruits (group)	1		
	Poultry meat	0.02	(*)	
	Rape seed	2		
	Raspberries, red, black	2		
	Rice, husked	0.1		
	Sugar cane	1		
	Tomato	1		
	Walnuts	0.05		
Thiacloprid		10		
		1		
		0.3		
		0.5		
		0.7		
		0.02	(*)	
		0.2		
		0.1		
		0.2		
		0.05		
		0.5		
		1		
		0.7		
		0.02	(*)	
		0.02	(*)	
		0.02	(*)	
		0.5		
		0.02	(*)	
		0.3		
		0.5		

		0.5		
		0.02		
		0.2		
		0.1		
		5		
		0.2		
		0.05		
		0.05		
		0.7		
		5		
		0.03	*	
		0.7		
		0.02		
	Dried Chilli	0.2		
Thiamethoxam	Artichoke, globe	0.5		
	Avocado	0.5		
	Banana	0.02	(*)	
	Barley	0.4		
	Barley, hay and/or straw	2		
	Beans with pods (Phaseolus spp.) immature pods and succulent seeds)	0.3		
	Berries and other small fruits	0.5		
	Brassica vegetables (except Brassica leafy vegetables) (group)	5		
	Cacao beans	0.02	(*)	
	Celery	1		
	Citrus fruits (group)	0.5		
	Coffee beans	0.2		
	Edible offal (mammalian)	0.01	(*)	
	Eggs	0.01	(*)	
	Fruiting vegetables, cucurbits (group)	0.5		
	Fruiting vegetables, other than cucurbits (group)	0.7		
	Hops, dry	0.09		
	Leafy vegetables	3		
	Legume vegetables (group)	0.01	(*)	
	Maize	0.05		
	Mango	0.2		

	Meat (from mammals other than marine mammals)	0.02		
	Milks	0.05		
	Mints	1.5		
	Oilseed	0.02	(*)	
	Papaya	0.01	(*)	
	Pecan	0.01	(*)	
	Peppers chili, dried	7		
	Pineapple	0.01	(*)	
	Pome fruits (group)	0.3		
	Popcorn	0.01	(*)	
	Poultry meat	0.01	(*)	
	Poultry, edible offal of	0.01	(*)	
	Pulses (group)	0.04		
	Root and tuber vegetables (group)	0.3		
	Stone fruits (group)	1		
	Sweet corn (corn-on-the-cob)	0.01	(*)	
	Tea, green, black (black, fermented and dried)	20		
	Wheat	0.05		
	Wheat, hay and/or straw	2		
	Rice	0.02		
	Cottonseed Oil	0.01		
	Tomato	0.7		
	Potato	0.3		
	Mustard seed	0.01		
	Cumin	0.01		
	Acid Lime	0.5		
	Ground nut	0.05	*	
	Ground nut oil	0.05	*	
	Sugarcane	0.05	*	
	Maize	0.05	*	
	Soybean	0.05	*	
	Soybean oil	0.05	*	
Thiifluzamide	Rice	0.05		
Thiodicarb	Cabbage	0.02		
	Brinjal	0.05		
	Red Gram	0.05		
	Black gram	0.03		
	Chilli	0.01		
	Dried Chilli	0.1		
	Cottonseed oil	0.02		

	Meats	0.02		
Thiophanate-Methyl	Apple	5		
	Papaya	7		
	Bottle gourd	0.4		
	Milks	0.05		
	Wheat	0.03	*	
	Pigeon pea	0.03	*	
	Cucumber	0.2		
	Grapes	3		
Tolfenpyrad	Bulb onions (subgroup)	0.09		
	Citrus oil, edible	80		
	Citrus pulp, dried	6		
	Edible offal (mammalian)	0.4		
	Eggs	0.01	(*)	
	Lemons and limes (including citron) (subgroup)	0.9		
	Mammalian fats (except milk fats)	0.01	(*)	
	Mandarins (including mandarin-like hybrids) (subgroup)	0.9		
	Meat (from mammals other than marine mammals)	0.01	(*)	
	Milks	0.01	(*)	
	Oranges, sweet, sour (including Orange-like hybrids) (subgroup)	0.6		
	Pecan	0.01	(*)	
	Peppers (subgroup)	0.5		(except martynia, okra and roselle)
	Peppers chili, dried	5		
	Potato	0.01	(*)	
	Poultry fats	0.01	(*)	
	Poultry meat	0.01	(*)	
	Poultry, edible offal of	0.01	(*)	
	Pummelo and grapefruits (including Shaddock-like hybrids, among others Grapefruit) (subgroup)	0.6		
	Tea, green	30		
	Cabbage	0.01	*	

	Okra	0.7		
Topramezone	Corn	0.05	*	
Triasulfuron	Wheat	0.01	*	
Triazophos	Cardamom	4		
	Coriander, seed	0.1		
	Cotton seed	0.2		
	Cotton seed oil, crude	1		
	Fennel, seed	0.1		
	Rice, polished	0.6		
	Soya bean (immature seeds)	0.5		
	Soya bean (young pod)	1		
	Spices, fruits and berries	0.07		(except cardamom)
	Spices, roots and rhizomes	0.1		
	Chilli	0.2		
	Dried Chilli	2		
Trichlorfon	Food grains	0.05		
	Milled food grains	0.01		
	Sugar beet	0.05		
	Fruits	0.1		
	Vegetables	0.1		
	Oil seeds	0.1		
	Edible oil (Refined)	0.05		
	Meat & Poultry	0.1		
	Milks	0.05		
Tricyclazole	Rice	3		
	Dried Chilli	3		
	Chilli	0.3		
Trifloxystrobin	Almond hulls	3		
	Asparagus	0.05	(*)	
	Banana	0.05		
	Barley	0.5		
	Barley, hay and/or straw	7		
	Beans (dry)	0.01	(*)	
	Brussels sprouts	0.1		
	Cabbages, head	1.5		
	Carrot	0.1		
	Celery	1		
	Citrus fruits (group)	0.5		
	Citrus pulp, dried	1		
	Cotton seed	0.4		
	Egg plant	0.7		

Eggs	0.04	(*)	
Flowerhead brassicas (includes Broccoli: Broccoli, Chinese and cauliflower)	0.5		
Fruiting vegetables, cucurbits (group)	0.3		
Ginseng	0.03	(*)	
Grape, dried (= Currants, Raisins and Sultanas)	5		
Grapes	3		
Hops, dry	40		
Kidney of cattle, goats, pigs and sheep	0.04	(*)	
Leek	0.7		
Lentil (dry)	0.01		
Lettuce, head	15		
Liver of cattle, goats, pigs & sheep	0.05		
Maize	0.02		
Meat (from mammals other than marine mammals)	0.05	(fat)	
Milks	0.02	(*)	
Olive oil, refined	1.2		
Olive oil, virgin	0.9		
Papaya	0.6		
Peanut	0.02	(*)	
Peas (dry)	0.01		
Peppers, sweet (including pimento or pimienta)	0.3		
Pome fruits (group)	0.7		
Potato	0.02	(*)	
Poultry meat	0.04	(*) (fat)	
Poultry, edible offal of	0.04	(*)	
Radish	0.08		
Radish leaves (including radish tops)	15		
Rice	5		
Rice bran, unprocessed	7		
Rice, hay and/or straw	10		
Soya bean (dry)	0.05		



	Spinach	20		
	Stone fruits (group)	3		
	Strawberry	1		
	Sugar beet	0.05		
	Sugar beet molasses	0.1		
	Sugar beet pulp, dry	0.2		
	Table olives	0.3		
	Tomato	0.7		
	Tree nuts (group)	0.02	(*)	
	Wheat	0.2		
	Wheat bran, unprocessed	0.5		
	Wheat, hay and/or straw	5		
	Chilly	0.4		
	Dry chilly	4		
	Mango	0.4		
	Apple	0.7		
	Banana	0.1		
	Maize	0.1		
	Cabbage	0.5		
Validamycin	Rice	0.01		
Zineb as CS2	Turmeric	2		
	Tea	0.1	*	

**Schedule-02**  
**Residue of Plant Growth Regulator**

Plant Growth Regulator	Names of the food	MRL (mg/kg)	Symbol	Notes
Ethephon	Milk	0.01	(*)	
	Apple	.8		
	Cherry, all commodities in this subgroup	5.00		
	Grapes	.8		
	Tomato	2.00		
	Cotton seed	6.00		
	Barley	1.5		
	Rye	.5		
	Wheat	.5		
	Cattle, goat and sheep meat	0.01	(*)	
	Poultry	0.08		
	Edible offal of poultry	0.20		
	Edible offal of cattle, goat and sheep	0.4		
	Eggs	0.01	(*)	
	Pineapple	2		
	Mangoes	2		
	Banana	2		

**Explanation for Symbol (s):**

(\*): At or about the limit of determination.<sup>[1]</sup>

\* Maximum Residue Limit fixed at Limit of Quantification (LOQ)<sup>[2]</sup>

**dry wt:** Dry Weight.

**Po:** The MRL accommodates post-harvest treatment of the commodity.

**(fat):** (for meat) The MRL/EMRL applies to the fat of meat.

**F:** Maximum Residue Limit Calculation on Fat Basis

**PoP:** (for processed foods) The MRL accommodates post-harvest treatment of the primary food commodity.

**Note:**

1. Limit of Detection (LOD): The lowest concentration or mass of the analyte that can be detected (but not quantified) in a sample. In practice, this is typically the analyte concentration at which the average signal/noise is 3.

2. Limit of quantification (LOQ): The smallest concentration of the analyte that can be quantified. It is commonly defined as the minimum concentration of the analyte in the test sample that can be determined with acceptable precision (repeatability) and accuracy under the stated conditions of the test. For the scope of this document, this is typically the analyte concentration at which the average signal/noise is 10.





**Schedule-3**  
**Portion of Commodity to Which the Maximum Residues Limits (MRL) Applies**

Classification of Commodities	Portion of Commodity to Which the MRL Applies (and Which is Analyzed)
<p><b>Root and Tuber Vegetables</b>  Root and tuber vegetables are starchy foods derived from the enlarged solid roots, tubers, corms or rhizomes, mostly subterranean, of various species of plants. The entire vegetable may be consumed.  Root and Tuber vegetables:  beets  radishes  carrots  sugar beets  celery  sweet potatoes  parsnips  turnips  potatoes  yams  radishes</p>	<p>Whole commodity after removing tops. Wash the roots or tubers in cold running water, brushing gently with a soft brush to remove loose soil and debris, if necessary, and then dab lightly with a clean tissue paper to dry. For the carrots, after drying, the tops are carefully cut off with a knife by cutting through the bottom of the stem at the lowest point of attachment of the outer petioles. If an annulus of root tissue is thereby severed from hollow-crown roots, the material should be recombined with the roots.</p>
<p><b>Bulb Vegetables</b>  Bulb vegetables are pungent flavourful foods derived from the fleshy scale bulbs, or growth buds of alliums of the lily family (Liliaceae). The entire bulb may be consumed following removal of the parchment like skin.  Bulb vegetables:  garlic  onions  leeks  spring onions</p>	<p>Bulb/dry onions and garlic:  Whole commodity after removal of roots and whatever parchment skin is easily detached.   Leeks and spring onions:  Whole vegetable after removal of roots and adhering soil.</p>
<p><b>Leafy Vegetables (Except Brassica Vegetables)</b>  Leafy vegetables (except Group 4 vegetables) are foods derived from the leaves of a wide variety of edible plants including leafy parts of Group 1 vegetables. The entire leaf may be consumed. Leafy vegetables of the brassica family are grouped separately.  Leafy vegetables:  beet leaves  radish leaves  corn salad  spinach  endive  sugar beet leaves  lettuce  Swiss chard</p>	<p>Whole commodity after removal of obviously decomposed or withered leaves.</p>
<p><b>Brassica (Cole) Leafy Vegetables</b>  Brassica (cole) leafy vegetables are foods derived from the leafy parts, stems and immature inflorescences of plants commonly known and botanically classified as brassicas and also known as cole vegetables. The entire vegetable may be consumed.</p>	<p>Whole commodity after removal of obviously decomposed or withered leaves. For cauliflower and headed broccoli analyse Brussels sprouts flower head and stems discarding leaves; for Brussels sprouts analyse "buttons" only.</p>




Brassica leafy vegetables: broccoli cauliflower Brussels sproutss collards cabbage kales cabbage, Chinese kohlrabi cabbage, red mustard greens cabbage, Savoy	
<b>Stem Vegetables</b> Stem vegetables are foods derived from the edible stems or shoots from a variety of plants. Stem vegetables: artichoke chicory (witloof) celery rhubarb	Whole commodity after removal of obviously decomposed or withered leaves.  Rhubarb and asparagus: stems only.  Celery and asparagus: remove adhering soil (e. rinsing in running water or by gentle brushing of dry commodity).
<b>Legume Vegetables</b> Legume vegetables are derived from the dried or succulent seeds and immature pods or leguminous plants commonly known as beans and peas. Succulent forms may be consumed as whole pods or as the shelled product. Legume fodder is in Group 18. Legume vegetables: beans navy beans broad beans runner beans dwarf beans snap beans French beans soybeans green beans peas kidney beans cow peas Lima beans sugar peas	Whole commodity.
<b>Fruiting Vegetables - Edible Peel</b> Fruiting vegetables - edible peel are derived from the immature or mature fruits of various plants, usually annual vines or bushes. The entire fruiting vegetables may be consumed. Fruiting vegetables - edible peel: cucumber pepper egg plant summer squash gherkin tomato okra	Whole commodity after removal of stems.
<b>Fruiting Vegetables - Inedible Peel</b>	Whole commodity after removal of stems.




<p>Fruiting vegetables - inedible peel are derived from the immature or mature fruits of various plants, usually annual vines or bushes. Edible portion is protected by skin, peel or husk which is removed or discarded before consumption.</p> <p>Fruiting vegetables - inedible peel:</p> <ul style="list-style-type: none"> <li>cantaloupe</li> <li>squash</li> <li>melon</li> <li>watermelon</li> <li>pumpkin</li> <li>winter squash</li> </ul>	
<p><b>Citrus Fruits</b></p> <p>Citrus fruits are produced by trees of the rue family and characterized by aromatic oily peels, globular form, and interior segments of juice filled vesicles. The fruit is fully exposed to pesticides during the growing season. The fruit pulp may be consumed in succulent form and as a beverage. The entire fruit may be used for preserving.</p> <p>Citrus fruits:</p>	Whole commodity.
<p><b>Pome Fruits</b></p> <p>Pome fruits are produced by trees related to the <i>genus pyrus</i> of the rose family (<i>Rosaceae</i>). They are characterized by fleshy tissue surrounding a core consisting of parchment like carpels enclosing the seed. The entire fruit, excepting the core, may be consumed in the succulent form or after processing.</p> <p>Pome fruits:</p> <ul style="list-style-type: none"> <li>apple</li> <li>quince</li> <li>pear</li> </ul>	Whole commodity after removal of stems.
<p><b>Stone Fruits</b></p> <p>Stone fruits are produced by trees related to the <i>genus prunus</i> of the rose family (<i>Rosaceae</i>) characterized by fleshy tissue surrounding a single hard shelled seed. The entire fruit, except seed, may be consumed in a succulent or processed form.</p> <p>Stone fruits:</p> <ul style="list-style-type: none"> <li>apricots</li> <li>nectarines</li> <li>cherries</li> <li>peaches</li> <li>sour cherries</li> <li>plums</li> <li>sweet cherries</li> </ul>	Whole commodity after removal of stems and stones but the residue calculated and expressed on the whole commodity without stem
<p><b>Small Fruits and Berries</b></p> <p>Small fruits and berries are derived from a variety of plants having fruit characterized by a high surface-weight ratio. The entire fruit, often including seed, may be consumed in a succulent or processed form.</p> <p>Small fruits and berries:</p> <ul style="list-style-type: none"> <li>blackberries</li> <li>gooseberries</li> <li>blueberries</li> <li>grapes</li> <li>boysenberries</li> <li>loganberries</li> </ul>	Whole commodity after removal of caps and stems. Currants: fruit with stems.


cranberries raspberries currants strawberries dewberries	
<b>Assorted Fruits - Edible Peel</b> Assorted fruits - edible peel are derived from the immature or mature fruits of a variety of plants, usually shrubs or trees from tropical or subtropical regions. The whole fruit may be consumed in a succulent or processed form. Assorted fruits - edible peel: dates olives figs	Dates and olives: whole commodity after removal of stems and stones but residue calculated and expressed on the whole fruit.  Figs: Whole commodity.
<b>Assorted Fruits - Inedible Peel</b> Assorted fruits - inedible peel are derived from the immature or mature fruits of different kinds of plants, usually shrubs or trees from tropical or subtropical regions. Edible portion is protected by skin, peel or husk. Fruit may be consumed in a fresh or processed form. Assorted fruits - inedible peel: avocados mangoes bananas papayas guavas passion fruits kiwi fruit pineapples	Whole commodity unless qualified.  Pineapples: after removal of crown.  Avocado and mangoes: whole commodity after removal of stone but calculated on whole fruit.  Bananas: after removal of crown tissues and stem
<b>Cereal Grains</b> Cereal grains are derived from the clusters of starchy seeds produced by a variety of plants primarily of the grass family ( <i>Gramineae</i> ). Husks are removed before consumption. Cereal grains: barley rye maize sorghum oats sweet corn rice wheat	Whole commodity.  Fresh corn and sweet corn: kernels plus cob with husk.
<b>Legume Oilseeds</b> Legume oilseeds are mature seeds from legumes cultivated for processing into edible vegetable oil or for direct use as human food. Legume oilseeds: peanuts	Whole kernel after removal of shell.
<b>Tree Nuts</b> Tree nuts are the seed of a variety of trees and shrubs which are characterized by a hard inedible shell enclosing an oil seed. The edible portion of the nut is consumed in succulent, dried or processed forms. Tree nuts: almonds macadamia nuts	Whole commodity after removal of shell.  Chestnuts: whole in skin.



chestnuts pecans filberts walnuts	
<b>Oilseed</b> Oilseed consists of the seed from a variety of plants used in the production of edible vegetable oils. Some important vegetable oilseeds are byproducts of fibre or fruit crops. Oilseed: cottonseed safflowerseed linseed sunflowerseed rapeseed	Whole commodity.
<b>Tropical Seeds</b> Tropical seeds consist of the seed from several tropical and semitropical trees and shrubs mostly used in the production of beverages and confections. Tropical seeds are consumed after processing. Tropical seeds: cacao beans coffee beans	Whole commodity.
<b>Herbs</b> Herbs consist of leaves, stems and roots from a variety of herbaceous plants used in relatively small amounts to flavour other foods. They are consumed in succulent or dried forms as components of other foods. Herbs:	Whole commodity.
<b>Spices</b> Spices consist of aromatic seeds, roots, fruits and berries from a variety of plants used in relatively small amount to flavour other foods. They are consumed primarily in the dried form as components of other foods. Spices:	Whole commodity.
<b>Teas</b> Teas are derived from the leaves of several plants, but principally <i>Camellia sinensis</i> . They are used in the preparation of infusions for consumption as stimulating beverages. They are consumed as extracts of the dried or processed product. Teas:	Whole commodity.
<b>Meats</b> Meats are the muscular tissue, including adhering fatty tissue from animal carcasses prepared for wholesale distribution. The entire product may be consumed. Meats: carcass meat (and carcass fat) carcass meat of cattle carcass meat of goats carcass meat of horses carcass meat of pigs carcass meat of sheep	Whole commodity. (For fat soluble pesticides a portion of carcass fat is analyzed and MRLs apply to carcass fat)
<b>Animal Fats</b>	Whole commodity.



<p>Animal fats are the rendered or extracted fat from the fatty tissue of animals. The entire product may be consumed.</p> <p>Animal fats:</p> <p>cattle fat</p> <p>sheep fat</p> <p>pig fat</p>	
<p><b>Meat Byproducts</b></p> <p>Meat byproducts are edible tissues and organs, other than meat and animal fat, from slaughtered animals as prepared for wholesale distribution.</p> <p>Examples: liver, kidney, tongue, heart. The entire product may be consumed.</p> <p>Meat byproducts (such as liver, kidney, etc.):</p> <p>cattle meat byproducts</p> <p>goat meat byproducts</p> <p>pig meat byproducts</p> <p>sheep meat byproducts</p>	Whole commodity.
<p><b>Milks</b></p> <p>Milks are the mammary secretion of various species of lactating herbivorous ruminant animals, usually domesticated. The entire product may be consumed.</p> <p>Milks:<sup>a</sup></p> <p>a - For milk and milk products regarding fat soluble pesticides see Codex Recommended Methods of Sampling for the Determination of Pesticide Residues for compliance with MRLs (CAC/GL 33-1999).</p>	Whole commodity.
<p><b>Milk Fats</b></p> <p>Milk fats are the rendered or extracted fats from milk.</p> <p>Milk fats:</p>	Whole commodity.
<p><b>Poultry Meats</b></p> <p>Poultry meats are the muscular tissues including adhering fat and skin from poultry carcasses as prepared for wholesale distribution. The entire product may be consumed.</p> <p>Poultry Meats:</p>	Whole commodity. (For fat soluble pesticides a portion of carcass fat is analyzed and MRLs apply to carcass fat).
<p><b>Poultry Fats</b></p> <p>Poultry fats are the rendered or extracted fats from fatty tissues of poultry. The entire product may be consumed.</p> <p>Poultry fats:</p>	Whole commodity.
<p><b>Poultry Byproducts</b></p> <p>Poultry byproducts are edible tissue and organs, other than poultry meat and poultry fat from slaughtered poultry.</p> <p>Poultry byproducts:</p>	Whole commodity.
<p><b>Eggs</b></p> <p>Eggs are the fresh edible portion of the reproductive body of several avian species. The edible portion includes egg white and egg yolk after removal of the shell.</p> <p>Eggs:</p>	Whole egg whites and yolks combined after removal of shells.

  
 By order of the Bangladesh Food Safety Authority  
**Md. Abdul Kayowm Sarker**  
 Chairman (Additional Secretary)